

## MODULES OVER GROUP RINGS OF LOCALLY FINITE GROUPS WITH FINITENESS RESTRICTIONS

OLGA DASHKOVA

ABSTRACT. We study an  $\mathbf{R}G$ -module  $A$ , where  $\mathbf{R}$  is a ring,  $A/C_A(G)$  is infinite,  $C_G(A) = 1$ ,  $G$  is a group. Let  $\mathfrak{L}_{\text{nf}}(G)$  be the system of all subgroups  $H \leq G$  such that the quotient modules  $A/C_A(H)$  are infinite. We investigate an  $\mathbf{R}G$ -module  $A$  such that  $\mathfrak{L}_{\text{nf}}(G)$  satisfies either the weak minimal condition or the weak maximal condition as an ordered set. It is proved that if  $G$  is a locally finite group then either  $G$  is a Chernikov group or  $G$  is a finite-finitary group of automorphisms of  $A$ .

### 1. INTRODUCTION

Important finiteness conditions in group theory are the weak minimal condition on subgroups and the weak maximal condition on subgroups. Let  $G$  be a group,  $\mathcal{M}$  be a set of subgroups of  $G$ .  $G$  is said to satisfy the weak minimal condition on  $\mathcal{M}$ -subgroups if for a descending series of subgroups  $G_0 \geq G_1 \geq G_2 \geq \cdots \geq G_n \geq G_{n+1} \geq \cdots$ ,  $G_n \in \mathcal{M}$ ,  $n \in \mathbb{N}$ , there exists the number  $m \in \mathbb{N}$  such that an index  $|G_n : G_{n+1}|$  is finite for any  $n \geq m$  [11]. Similarly  $G$  is said to satisfy the weak maximal condition on  $\mathcal{M}$ -subgroups if for an ascending series of subgroups  $G_0 \leq G_1 \leq G_2 \leq \cdots \leq G_n \leq G_{n+1} \leq \cdots$ ,  $G_n \in \mathcal{M}$ ,  $n \in \mathbb{N}$ , there exists the number  $m \in \mathbb{N}$  such that an index  $|G_n : G_{n+1}|$  is finite for any  $n \geq m$  [1].

These finiteness conditions were applied to investigate infinite dimensional linear periodic groups [9]. Also similar finiteness conditions were considered in [2].

Let  $A$  be an  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring,  $G$  be a group.  $G$  is a finite-finitary group of automorphisms of  $A$  if  $C_G(A) = 1$  and  $A/C_A(g)$  is finite for any  $g \in G$  [10]. Finite-finitary groups of automorphisms of  $A$  with additional restrictions were studied in [10].

---

2010 *Mathematics Subject Classification.* 20F19, 20H25.

*Key words and phrases.* group ring, locally finite group, locally solvable group.

Let  $\mathfrak{L}_{\text{nf}}(G)$  be the system of all subgroups  $H$  of  $G$  such that  $A/C_A(H)$  is infinite. Previously, we studied an  $\mathbf{R}G$ -module  $A$  with some restrictions on subgroups of  $\mathfrak{L}_{\text{nf}}(G)$  [4, 3, 5].

In this paper we continue these investigations. We say that  $G$  satisfies the condition  $W_{\text{min-nf}}$  if  $G$  satisfies the weak minimal condition on  $\mathcal{M}$ -subgroups where  $\mathcal{M} = \mathfrak{L}_{\text{nf}}(G)$  and  $G$  satisfies the condition  $W_{\text{max-nf}}$  if  $G$  satisfies the weak maximal condition on  $\mathcal{M}$ -subgroups where  $\mathcal{M} = \mathfrak{L}_{\text{nf}}(G)$ .

Next, we consider an  $\mathbf{R}G$ -module  $A$  with  $C_G(A) = 1$ . We investigate an  $\mathbf{R}G$ -module  $A$  such that  $G$  satisfies either  $W_{\text{min-nf}}$  or  $W_{\text{max-nf}}$ . Main results of the paper are Theorem 1 and Theorem 2.

## 2. PRELIMINARY RESULTS

**Lemma 1.** *Let  $A$  be an  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring. Then the following conditions hold:*

- (1) *if  $L \leq H \leq G$  and  $A/C_A(H)$  is finite then  $A/C_A(L)$  is finite also;*
- (2) *if  $L, H \leq G$ ,  $A/C_A(L)$  and  $A/C_A(H)$  are finite then  $A/C_A(\langle L, H \rangle)$  is finite also.*

**Corollary 1.** *Let  $A$  be an  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring,  $FFD(G)$  be the set of all elements  $x \in G$  such that  $A/C_A(x)$  is finite. Then  $FFD(G)$  is a normal subgroup of  $G$ .*

*Proof.* By Lemma 1 (2)  $FFD(G)$  is a subgroup of  $G$ . Since  $C_A(x^g) = C_A(x)g$  for each  $x, g \in G$  then  $FFD(G)$  is a normal subgroup of  $G$ .  $\square$

**Lemma 2.** *Let  $A$  be an  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring,  $H$  be a subgroup of  $G$ . Suppose that  $H$  contains a normal subgroup  $K$  such that  $A/C_A(K)$  is infinite. Then the following conditions hold:*

- (1) *if  $G$  satisfies  $W_{\text{min-nf}}$  then  $H/K$  satisfies the weak condition of minimality on subgroups;*
- (2) *if  $G$  satisfies  $W_{\text{max-nf}}$  then  $H/K$  satisfies the weak condition of maximality on subgroups.*

**Lemma 3.** *Let  $A$  be an  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring,  $L$ ,  $K$  and  $H$  be subgroups of  $G$  with the following properties:*

- (i)  *$K$  is a normal subgroup of  $L$ ;*
- (ii)  *$K$  and  $L$  are  $H$ -invariant subgroups;*
- (iii)  *$L/K \cap HK/K = \langle 1 \rangle$ ;*
- (iv)  *$L/K = \text{Dr}_{n \in \mathbb{N}} L_n/K$ ,  $L_n/K \neq \langle 1 \rangle$  is an  $H$ -invariant subgroup for any  $n \in \mathbb{N}$ .*

*Then the following conditions hold:*

- (1) *if  $G$  satisfies  $W_{\text{max-nf}}$  then  $A/C_A(HL)$  is finite;*
- (2) *if  $G$  satisfies  $W_{\text{min-nf}}$  then  $A/C_A(HK)$  is finite.*

*Proof.* There are two infinite subsets  $\Sigma$  and  $\Delta$  of  $\mathbb{N}$  such that  $\Sigma \cup \Delta = \mathbb{N}$ ,  $\Sigma \cap \Delta = \emptyset$ . Since  $\Delta$  is infinite then there is an infinite strongly ascending

series of subsets of  $\Delta$

$$\Delta(1) \subset \Delta(2) \subset \cdots \subset \Delta(k) \subset \cdots .$$

Also there is strongly descending series of subsets of  $\Delta$

$$\Delta^*(1) \supset \Delta^*(2) \supset \cdots \supset \Delta^*(k) \supset \cdots ,$$

such that the sets  $\Delta(k+1) \setminus \Delta(k)$  and  $\Delta^*(k) \setminus \Delta^*(k+1)$  are infinite for any  $n \in \mathbb{N}$ . Let

$$D_k/K = Dr_{t \in \Sigma \cup \Delta(k)} L_t/K$$

and

$$D_k^*/K = Dr_{t \in \Sigma \cup \Delta^*(k)} L_t/K.$$

At first we consider the strongly ascending series of subgroups

$$HD_1 < HD_2 < \cdots < HD_k < \cdots .$$

$|HD_{k+1} : HD_k|$  are infinite by construction. If  $G$  satisfies  $W_{\max\text{-nf}}$  then there is  $m \in \mathbb{N}$  such that  $A/C_A(HD_m)$  is finite. Since  $\langle H, L_t | t \in \Sigma \rangle \leq HD_m$  then  $A/C_A(\langle H, L_t | t \in \Sigma \rangle)$  is finite by Lemma 1. Similarly we prove that  $A/C_A(\langle H, L_t | t \in \Delta \rangle)$  is finite.

Since  $\Sigma \cup \Delta = \mathbb{N}$  we obtain

$$\langle \langle H, L_t | t \in \Delta \rangle, \langle H, L_t | t \in \Sigma \rangle \rangle = \langle H, L_t | t \in \Sigma \cup \Delta \rangle = HL.$$

By Lemma 1  $A/C_A(HL)$  is finite.

Likewise we can construct the strongly descending series of subgroups

$$HD_1^* > HD_2^* > \cdots > HD_k^* > \cdots ,$$

such that  $|HD_k^* : HD_{k+1}^*|$  are infinite. If  $G$  satisfies  $W_{\min\text{-nf}}$  then there is  $m \in \mathbb{N}$  such that  $A/C_A(HD_m^*)$  is finite. Since  $HK \leq HD_m^*$  then  $A/C_A(HK)$  is finite by Lemma 1.  $\square$

**Corollary 2.** *Let  $A$  be an  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring,  $L, K$  and  $H$  be subgroups of  $G$  with the the following properties:*

- (i)  $K$  is a normal subgroup of  $L$ ;
- (ii)  $K$  and  $L$  are  $H$ -invariant subgroups;
- (iii)  $L/K = Dr_{n \in \mathbb{N}} L_n/K$  where  $L_n/K \neq \langle 1 \rangle$  is an  $H$ -invariant subgroup for any  $n \in \mathbb{N}$ ;
- (iv) the set  $\mathbb{N} \setminus \text{Supp}(L/K \cap HK/K)$  is infinite.

*If  $G$  satisfies either  $W_{\min\text{-nf}}$  or  $W_{\max\text{-nf}}$  then  $A/C_A(HK)$  is finite. In particular  $A/C_A(H)$  is finite.*

*Proof.* Let  $\Delta = \mathbb{N} \setminus \text{Supp}(L/K \cap HK/K)$  and  $T/K = Dr_{n \in \Delta} L_n/K$ . Then  $T/K \cap HK/K = \langle 1 \rangle$ . We apply Lemma 3.  $\square$

**Corollary 3.** *Let  $A$  be an  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring,  $L, K$  and  $H$  be subgroups of  $G$  with the the following properties:*

- (i)  $K$  is a normal subgroup of  $L$ ;
- (ii)  $K$  and  $L$  are  $H$ -invariant subgroups;

(iii)  $L/K = Dr_{n \in \mathbb{N}} L_n/K$ ,  $L_n/K \neq \langle 1 \rangle$  is an  $H$ -invariant subgroup for any  $n \in \mathbb{N}$ .

If  $G$  satisfies either  $W_{\min\text{-nf}}$  or  $W_{\max\text{-nf}}$  then  $A/C_A(\langle h \rangle K)$  is finite for any  $h \in H$ . In particular  $H \leq \text{FFD}(G)$ .

*Proof.* Let  $h \in H$ . Since  $L_n/K$  is an  $H$ -invariant subgroup for any  $n \in \mathbb{N}$  then  $L_n/K$  is an  $\langle h \rangle$ -invariant subgroup for any  $n \in \mathbb{N}$ . In particular the set  $\text{Supp}(\langle h \rangle K/K \cap L/K)$  is finite. Then  $A/C_A(\langle h \rangle K)$  is finite by Corollary 2.  $\square$

### 3. MAIN RESULTS

Obviously that a Chernikov group satisfies both the weak minimal condition on subgroups and the weak maximal condition on subgroups. It follows that if  $A$  is an  $\mathbf{R}G$ -module and  $G$  is Chernikov then  $G$  satisfies both  $W_{\min\text{-nf}}$  and  $W_{\max\text{-nf}}$ .

**Lemma 4.** *Let  $A$  be an  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring. Suppose that  $G$  satisfies either  $W_{\min\text{-nf}}$  or  $W_{\max\text{-nf}}$ . Let  $K$  and  $H$  be subgroups of  $G$  such that  $K$  is a normal subgroup of  $H$  and  $H/K$  is an infinite elementary abelian  $p$ -group for some prime  $p$ . Suppose that  $K$  and  $H$  are  $\langle g \rangle$ -invariant for some  $g \in G$ . If  $g^k \in C_G(H/K)$  for some  $k \in \mathbb{N}$  then  $g \in \text{FFD}(G)$ .*

*Proof.* Let  $M = H/K$ . We take  $1 \neq b_1 \in M$ . Put  $B_1 = \langle b_1 \rangle^{\langle g \rangle}$ . Since  $g$  induces the automorphism of finite order on  $M$  then  $B_1$  is finite.  $M = B_1 \times C_1$  is valid for some subgroup  $C_1$ .

Let

$$\{C_1^y | y \in \langle g \rangle\} = \{U_1, \dots, U_m\}.$$

It follows that the  $\langle g \rangle$ -invariant subgroup

$$D_1 = U_1 \cap \dots \cap U_m = \text{Core}_{\langle g \rangle}(C_1)$$

has finite index in  $M$ . Let  $1 \neq b_2 \in D_1$  and  $B_2 = \langle b_2 \rangle^{\langle g \rangle}$ . Then  $\langle B_1, B_2 \rangle = B_1 \times B_2$ . As before we conclude that  $M = (B_1 \times B_2) \times C_2$  for some subgroup  $C_2$ . Similarly we can construct the infinite set  $\{B_n | n \in \mathbb{N}\}$  of non-trivial  $\langle g \rangle$ -invariant subgroups such that  $\langle B_n | n \in \mathbb{N} \rangle = Dr_{n \in \mathbb{N}} B_n$ . By Corollary 3 we have that  $g \in \text{FFD}(G)$ .  $\square$

Let  $\pi(G)$  be the set of all prime divisors of orders of elements of  $G$ .

**Corollary 4.** *Let  $A$  be an  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring. Suppose that  $G$  satisfies either  $W_{\min\text{-nf}}$  or  $W_{\max\text{-nf}}$ . Let  $K$  and  $H$  are subgroups of  $G$  such that  $K$  is a normal subgroup of  $H$ ,  $H/K$  is a periodic almost locally solvable group. If  $H/K$  is not Chernikov then  $H \leq \text{FFD}(G)$ .*

*Proof.* Let  $L/K$  be a locally solvable normal subgroup of  $H/K$  of finite index. Since  $H/K$  is not Chernikov then  $L/K$  is not Chernikov too. Let  $g$  be an element of  $H$ . Then  $L/K$  contains an abelian  $\langle g \rangle$ -subgroup  $C/K$  which is not Chernikov [12]. If the set  $\pi(C/K)$  is infinite then  $g \in \text{FFD}(G)$  by Corollary

3. If  $\pi(C/K)$  is finite then there is the prime  $p$  such that Sylow  $p$ -subgroup  $P/K$  of  $C/K$  is not Chernikov. It follows that the lower layer  $B/K$  of  $P/K$  is infinite. Therefore  $L/K$  contains a  $\langle g \rangle$ -invariant infinite elementary abelian subgroup  $B_1/K$ . Then  $g \in FFD(G)$  by Lemma 4.  $\square$

**Corollary 5.** *Let  $A$  be an  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring. Suppose that  $G$  satisfies either  $W_{min-nf}$  or  $W_{max-nf}$ . Let  $K$  and  $H$  be subgroups of  $G$  such that  $K$  is a normal subgroup of  $H$ ,  $H/K$  is a locally finite group. If  $H/K$  is not Chernikov then  $H \leq FFD(G)$ .*

*Proof.* Let  $g$  be an element of  $H$  and  $C/K = C_{H/K}(gK)$ . If  $C/K$  is not Chernikov then by Theorem [8]  $C/K$  contains an abelian subgroup  $D/K$  which is a direct product of infinite set of non-trivial cyclic subgroups. By Corollary 3 we have that  $g \in FFD(G)$ . Suppose that  $C/K$  is not Chernikov. Then  $H/K$  is an almost locally solvable group [6] and  $g \in FFD(G)$  by Corollary 4. We have that  $H \leq FFD(G)$ .  $\square$

It follows that Theorem 1 is valid.

**Theorem 1.** *Let  $A$  be an  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring,  $G$  be a locally finite group. If  $G$  satisfies either  $W_{min-nf}$  or  $W_{max-nf}$  then either  $G$  is a Chernikov group or  $G$  is a finite-finitary group of automorphisms of  $A$ .*

**Lemma 5.** *Let  $A$  be a  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring,  $G$  be a locally solvable group. Suppose that  $A/C_A(G)$  is finite. Then  $G$  is almost abelian.*

*Proof.* Let  $C = C_A(G)$ . Then  $A$  has the series of  $\mathbf{R}G$ -submodules  $\langle 0 \rangle \leq C \leq A$ , where  $A/C$  is a finite  $\mathbf{R}$ -module. Since  $G \leq C_G(C)$  then  $G/C_G(C)$  is trivial. As  $A/C$  is a finite  $\mathbf{R}$ -module then  $G/C_G(A/C)$  is finite.

Let  $H = C_G(C) \cap C_G(A/C)$ . Each element of  $H$  acts trivially on every factor of the series  $\langle 0 \rangle \leq C \leq A/C$ . By Kaluzhnin Theorem (p. 144 [7])  $H$  is abelian. By Remak's Theorem

$$G/H \leq G/C_G(C) \times G/C_G(A/C).$$

It follows that  $G/H$  is finite. Then  $G$  is an almost abelian group.  $\square$

Let  $G_{\mathfrak{S}}$  be the intersection of all normal subgroups  $K$  of  $G$  such that  $G/K$  is solvable. If  $G$  is a solvable group then we denote the step of solvability of  $G$  by  $s(G)$ .

**Theorem 2.** *Let  $A$  be an  $\mathbf{R}G$ -module,  $\mathbf{R}$  be an associative ring,  $G$  be a locally solvable periodic group. If  $G$  satisfies either  $W_{min-nf}$  or  $W_{max-nf}$  then  $G/G_{\mathfrak{S}}$  is solvable.*

*Proof.* Otherwise  $H = G/G_{\mathfrak{S}}$  is unsolvable. Let  $F_1$  be a finite subgroup of  $H$ . Since  $H$  is approximated by solvable subgroups then there is a normal subgroup  $K_1$  of  $H$  such that  $F_1 \cap K_1 = \langle 1 \rangle$  and  $H/K_1$  is solvable. It follows that  $K_1$  is unsolvable. Therefore the steps of solvability of finite subgroups of  $K_1$  not limited by the number. Then  $K_1$  contains a finite subgroup  $D_1$  such

that  $s(F_1) < s(D_1)$ . Since  $F_1$  and  $D_1$  are finite then they are solvable. Let  $F_2 = D_1^{F_1}$ . Then  $F_2$  is a finite  $F_1$ -invariant subgroup such that  $s(F_1) < s(F_2)$ . Since  $F_1F_2$  is finite there is a normal subgroup  $K_2$  of  $H$  such that  $F_1F_2 \cap K_2 = \langle 1 \rangle$  and  $H/K_2$  is solvable. Since  $K_2$  is unsolvable then we can choose a finite  $F_1F_2$ -invariant subgroup  $F_3$  of  $K_2$  such that  $s(F_2) < s(F_3)$ . Continuing our reasoning, we construct the strongly ascending series of finite subgroups  $F_1 < F_1F_2 < \dots < F_1F_2 \dots F_n < \dots$  with the the following properties:

- (i)  $F_n$  is an  $F_j$ -invariant subgroup for  $j < n$ ;
- (ii)  $s(F_j) < s(F_n)$  for  $j < n$ ;
- (iii)  $F_1F_2 \dots F_n \cap \langle F_j | j > n \rangle = \langle 1 \rangle$  for any  $n \in \mathbb{N}$ .

It follows that  $\langle F_j | j \in \Delta \rangle$  is decomposed in the direct product of  $F_j, j \in \Delta$ , for an infinite subset  $\Delta$  of  $\mathbb{N}$ . Therefore  $\langle F_j | j \in \Delta \rangle$  is unsolvable.

At first we suppose that  $G$  satisfies  $W_{\min\text{-nf}}$ . There is an infinite strictly descending series of subsets

$$\mathbb{N} \supset \Delta(1) \supset \Delta(1) \supset \dots \supset \Delta(k) \supset \dots$$

such that  $\Delta(k) \setminus \Delta(k+1)$  is infinite for any  $k \in \mathbb{N}$ . Let  $L_k = \langle F_j | j \in \Delta(k) \rangle$  for any  $k \in \mathbb{N}$ . We obtain the strongly descending series of subgroups  $L_1 > L_2 > \dots > L_k > \dots$  of  $H$ . Let  $M_k$  is the preimage of  $L_k$  in  $G$ . Then  $M_1 > M_2 > \dots > M_k > \dots$  is the strongly descending series of subgroups of  $G$  such that  $|M_k : M_{k+1}|$  are infinite. Hence there is  $t \in \mathbb{N}$  such that  $A/C_A(M_t)$  is finite.  $M_t$  is solvable by Lemma 5. It follows that  $L_t = M_t/G_{\mathfrak{E}}$  is solvable. Previously, we proved that  $L_t = \langle F_j | j \in \Delta(t) \rangle$  is unsolvable. Contradiction.

If  $G$  satisfies  $W_{\max\text{-nf}}$  we construct an infinite strictly ascending series of subsets of  $\mathbb{N}$  and conduct similar reasoning.  $\square$

When writing the paper the author used the methods of [9].

## REFERENCES

- [1] R. Baer. Polyminimaxgruppen. *Math. Ann.*, 175:1–43, 1968.
- [2] V. A. Bovdi and V. P. Rudko. Indecomposable linear groups. *Indian J. Pure Appl. Math.*, 40:327–336, 2009.
- [3] O. Dashkova. Modules over group rings of locally soluble groups with a certain condition of minimality. *Miskolc Math. Notes*, 15:383–392, 2014.
- [4] O. Y. Dashkova. On one class of modules over group rings with finiteness restrictions. *Int. J. Group Theory*, 3:37–46, 2014.
- [5] O. Y. Dashkova. Modules over group rings of groups with restrictions on the system of all proper subgroups. *Int. J. Group Theory*, 4:43–48, 2015.
- [6] B. Hartley. Fixed points of automorphisms of certain locally finite groups and chevalley groups. *J. London Math. Soc.*, 37:421–436, 1988.
- [7] M. I. Kargapolov and Y. I. Merzlyakov. *Osnovy teorii grupp*. Izdat. “Nauka”, Moscow, 1972.
- [8] O. Kegel and B. Wehrfritz. *Locally Finite Groups*. North-Holland Mathematical Library. North-Holland, Amsterdam, London, New York, 1973.

- [9] J. M. Munoz-Escolano, J. Otal, and N. N. Semko. Periodic linear groups with the weak chain conditions on subgroups of infinite central dimension. *Comm. Algebra*, 36:749–763, 2008.
- [10] B. Wehrfritz. Finite-finitary groups of automorphisms. *J. Algebra Appl.*, 1:375–389, 2002.
- [11] D. I. Zaitzev. Groups satisfying the the weak minimal condition. *Ukr. Math. J.*, 20:408–416, 1968.
- [12] D. I. Zaitzev. On solvable subgroups of locally soluble groups. *Rep. USSR Academy of Sciences*, 214:1250–1253, 1974.

*Received May 2, 2016.*

THE BRANCH OF MOSCOW STATE UNIVERSITY IN SEVASTOPOL,  
HEROES OF SEVASTOPOL STREET 7,  
SEVASTOPOL, 299001, RUSSIA  
*E-mail address:* odashkova@yandex.ru