

Molecular, morphological and acoustic assessment of the genus *Ophryophryne* (Anura, Megophryidae) from Langbian Plateau, southern Vietnam, with description of a new species

Nikolay A. Poyarkov Jr.^{1,2}, Tang Van Duong^{1,3}, Nikolai L. Orlov⁴,
Svetlana S. Gogoleva^{2,5,6}, Anna B. Vassilieva^{1,2,5}, Luan Thanh Nguyen⁷,
Vu Dang Hoang Nguyen⁷, Sang Ngoc Nguyen⁷, Jing Che^{8,9}, Stephen Mahony^{10,11}

1 Department of Vertebrate Zoology, Biological faculty, Lomonosov Moscow State University, Leninskiye Gory, Moscow, GSP-1, 119991, Russia **2** Joint Russian-Vietnamese Tropical Research and Technological Center, 63 Nguyen Van Huyen Road, Nghia Do, Cau Giay, Hanoi, Vietnam **3** Vietnam National Museum of Nature, Vietnam Academy of Science and Technology (VAST), 18 Hoang Quoc Viet Road, Hanoi, Vietnam **4** Zoological Institute, Russian Academy of Sciences, Universitetskaya nab., 1, St. Petersburg, 199034, Russia **5** A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Leninskii pr., 33, Moscow 119071, Russia **6** Zoological Museum of the Lomonosov Moscow State University, Bolshaya Nikiiskaya st. 6, Moscow 125009, Russia **7** Institute of Tropical Biology, Vietnam Academy of Science and Technology (VAST), 85 Tran Quoc Toan St., District 3, Ho Chi Minh City, Vietnam **8** State Key Laboratory of Genetic Resources and Evolution State, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650223, Yunnan, China **9** Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences, Yezin, Nay Pyi Taw 05282, Myanmar **10** UCD School of Biology and Environmental Science, UCD Science Centre (West), University College Dublin, Belfield, Dublin 4, Ireland **11** Department of Life Sciences, The Natural History Museum, South Kensington, London, United Kingdom

Corresponding authors: Nikolay A. Poyarkov (n.poyarkov@gmail.com); Stephen Mahony (stephenmahony2@gmail.com)

Academic editor: A. Crottini | Received 23 September 2016 | Accepted 20 March 2017 | Published 3 May 2017

<http://zoobank.org/DFAF2079-2883-4BA9-931E-07F00D1F7D76>

Citation: Poyarkov Jr NA, Duong TV, Orlov NL, Gogoleva SS, Vassilieva AB, Nguyen LT, Nguyen VDH, Nguyen SN, Che J, Mahony S (2017) Molecular, morphological and acoustic assessment of the genus *Ophryophryne* (Anura, Megophryidae) from Langbian Plateau, southern Vietnam, with description of a new species. ZooKeys 672: 49–120. <https://doi.org/10.3897/zookeys.672.10624>

Abstract

Asian Mountain Toads (*Ophryophryne*) are a poorly known genus of mostly small-sized anurans from southeastern China and Indochina. To shed light on the systematics within this group, the most com-

prehensive mitochondrial DNA phylogeny for the genus to date is presented, and the taxonomy and biogeography of this group is discussed. Complimented with extensive morphological data (including associated statistical analyses), molecular data indicates that the Langbian Plateau, in the southern Annamite Mountains, Vietnam, is one of the diversity centres of this genus where three often sympatric species of *Ophryophryne* are found, *O. gerti*, *O. synoria* and an undescribed species. To help resolve outstanding taxonomic confusion evident in literature (reviewed herein), an expanded redescription of *O. gerti* is provided based on the examination of type material, and the distributions of both *O. gerti* and *O. synoria* are considerably revised based on new locality records. We provide the first descriptions of male mating calls for all three species, permitting a detailed bioacoustics comparison of the species. We describe the new species from highlands of the northern and eastern Langbian Plateau, and distinguish it from its congeners by a combination of morphological, molecular and acoustic characters. The new species represents one of the smallest known members of the genus *Ophryophryne*. At present, the new species is known from montane evergreen forest between 700–2200 m a.s.l. We suggest the species should be considered Data Deficient following IUCN's Red List categories.

Keywords

12S rRNA, 16S rRNA, advertisement call, amphibian, biodiversity, Da Lat Plateau, frog, Indochina, southeast Asia, taxonomy, Truong Son

Introduction

Asian Mountain toads (*Ophryophryne* Boulenger, 1903) are a small group of frogs from southeast Asia with a rather limited distribution mostly in mountains of eastern Indochina and adjacent parts of southern China (Yunnan and Guangxi) and northern Thailand (Inger et al. 1999, Ohler 2003, Orlov and Ananjeva 2007, Yang 2008). The genus *Ophryophryne* is still poorly studied, to date five (Ohler 2003, Stuart et al. 2006) or six (Stuart et al. 2010) species are recognized, with little consensus on the taxonomic status of several forms. All of the known *Ophryophryne* species have been reported from the Truong Son or Annamite mountains in Vietnam, which may be considered as an area of highest diversity for this group (Orlov and Ananjeva 2007).

The systematic status of *Ophryophryne* has long been a source of confusion. Boulenger (1903) described the genus and species *O. microstoma* Boulenger, 1903, and though he clearly stated that *Ophryophryne* is closely allied to *Megophrys* Kuhl & Van Hasselt, 1822, he noted characters also shared by members of the family Bufonidae Gray, 1825 (lacking vomerine and maxillary teeth and presence of horizontal pupil). Subsequently, Noble (1926), mostly based on morphology of the pectoral girdle, clearly demonstrated that the genus *Ophryophryne* is a member of Pelobatidae (at the time including the subfamily Megophryinae Bonaparte, 1850), and assumed its close affinities to *Megophrys*. However, due to widespread misinterpretation of Boulenger's original statement, the genus *Ophryophryne* was nevertheless incorrectly listed as a member of Bufonidae in several classical works on batrachians (Bourret 1937, 1942, Gorham 1974, Guibé 1950, Nguyen and Ho 1996, Taylor 1962).

The systematic status of the genus *Ophryophryne* among the Megophryidae has been discussed in several works. Liu and Hu (1962) provided the first description of the *Ophryophryne* tadpole which was remarkably similar to those of *Megophrys*, which led Dubois (1980) to rank *Ophryophryne* at the level of subgenus within *Megophrys*. Soon afterwards, Dubois re-evaluated his proposition and elevated *Ophryophryne* back to the genus-level status (Dubois 1987). Summarizing available cytological, morphological and ecological evidence, Rao and Yang (1997) proposed to split *Megophrys s. lato*, regarding *Ophryophryne* as a separate genus, as well as the former *Megophrys* subgenera, *Megophrys s. stricto*, *Atympanophrys* Tian & Hu, 1983, *Brachytarsophrys* Tian & Hu, 1983 and *Xenophrys* Günther, 1864. Several studies indicated close affinities of *Ophryophryne* to the genus *Xenophrys* (Tian and Hu 1983, 1985, Frost 1985, Ye et al. 1993, Rao and Yang 1997, Manthey and Grossmann 1997). Delorme et al. (2006) recognized the tribe Xenophryini, containing two genera *Ophryophryne* and *Xenophrys*, and most recent faunal reviews have treated *Ophryophryne* as a valid genus within Megophryidae (Fei et al. 1999, 2005, 2009, 2010, 2012, Orlov and Ananjeva 2007, Nguyen et al. 2009). Recently, Mahony (2011a) suggested that insufficient evidence was available for the morphological distinction of *Xenophrys* from *Megophrys* and suggested to retain the historical usage of *Megophrys s. lato* for species of both genera pending a taxonomic review of the group, however, he did not discuss the status of *Ophryophryne*.

Though a comprehensive phylogeny of the genus *Ophryophryne* is still pending, preliminary molecular data were contradictory, suggesting both as sister-clade relationships of *Ophryophryne* with respect to a monophyletic group composed of *Xenophrys*, *Megophrys*, and *Brachytarsophrys* (Pyron and Wiens 2011), or providing evidence of the paraphyly of *Xenophrys* with respect to *Ophryophryne* (e.g. Wang et al. 2012). A recent phylogenetic study on Megophryinae by Chen et al. (2017) provides new insights on evolutionary relationships within this group, indicating contrasting (though poorly supported) phylogenetic positions of *Ophryophryne* in their multilocus nuclear-gene based phylogeny and matrilineal mtDNA genealogy. Chen et al. (2017) preliminarily recognized *Ophryophryne* as one of the five monophyletic genera within Megophryinae (*Ophryophryne*, *Brachytarsophrys*, *Xenophrys*, *Atympanophrys* and *Megophrys*). However, Mahony et al. (2017) provides an alternative hypothesis based on extensive morphological studies and a larger nuclear gene dataset. They provided compelling evidence (recent diversification, insufficient morphological or biological distinction of major clades) for the consideration of Megophryinae to be treated as a single genus, *Megophrys*, with seven sub-clades (including *Ophryophryne*) being treated as subgenus level taxa. Their phylogenetic analyses provided strong support for the sister taxa relationship of *Ophryophryne* and a clade corresponding to *Panophrys* (previously considered a synonym or subclade of *Xenophrys*, e.g., Delorme et al. 2006, Chen et al. 2017).

For a long time after its' description, the genus *Ophryophryne* was thought to include a single species, *O. microstoma*, described from "Mau Son" in Tonkin (northern Vietnam). Later, Bourret (1937) described a second species, *O. poilani* Bourret, 1937, based upon a single, badly preserved specimen from "Dong Tam Ve" in Quang Tri Prov. of Annam (central Vietnam). Almost half a century later a third species, *O. pachy-*

proctus Kou, 1985, was described by Kou (1985) from Mengla County in Yunnan Prov. (southern China). Ohler (2003) revised the available material on the genus and, mostly based on samples collected by M. Smith (in southern Vietnam), and I.S. Darevsky and N.L. Orlov (in central Vietnam), described two more species: *O. gerti* Ohler, 2003 and *O. hansi* Ohler, 2003, respectively (the type series of *O. gerti* included specimens from the Langbian Plateau in southern Vietnam, and Laos). Ohler (2003) revised diagnostic characters and provided a key for the genus; she also examined the type specimen of *O. poilani* and considered it to be a junior synonym of *O. microstoma* (opinion not shared by Stuart et al. 2010). The last major progress on the taxonomy of *Ophryophryne* was made by Stuart et al. (2006), who reported the genus for Cambodia and described one more species, *O. synoria* Stuart, Sok & Neang, 2006 from Mondolkiri Prov. in eastern hilly Cambodia, near the Vietnamese border. A recent review of southern Vietnamese herpetofauna by Vassilieva et al. (2016) based on morphological evidence recorded *O. synoria* for lowland areas of Dong Nai and Binh Phuoc provinces.

The Langbian (or Da Lat) Plateau forms the southernmost edge of the Annamite Mountains, or Truong Son Range, a mountain chain spanning the breadth of Indochina, including parts of Vietnam, Laos and Cambodia. To date, following the review by Ohler (2003), only *Ophryophryne gerti* has been recorded from the high elevations (above 1000 m a.s.l.) of the Langbian Plateau (Stuart et al. 2010, Nguyen et al. 2009, 2014). However, our recent fieldwork in this area from 2007 until 2016 revealed the presence of at least three morphologically distinct species, often recorded in syntopy (Orlov et al. 2008, Poyarkov and Vassilieva 2011). Further investigation of partial 12S rRNA–16S rRNA mtDNA gene sequences, as well as the study of advertisement calls from the Langbian *Ophryophryne* populations, herein confirm their specific status and reveals that one of the lineages represents a previously undescribed species. We also provide the first preliminary mtDNA phylogeny for the genus *Ophryophryne* and discuss the biogeography of the genus in Indochina in light of our new data.

Materials and methods

Sample collection. All specimens were collected during fieldwork in southern Vietnam in 2007–2016. Frogs were collected mostly during night excursions by opportunistic visual searching, or by sound when calling. Geographic coordinates were obtained using a Garmin GPSMAP 60CSx GPS receiver and recorded in datum WGS 84. The geographic position of the surveyed localities and the distribution of *Ophryophryne* species in the southern Annamite Mountains (Truong Son) and adjacent regions of southern Indochina (eastern Cambodia) are shown in Fig. 1. The newly collected specimens were deposited in the herpetological collection of the Zoological Museum of Moscow State University in Moscow, Russia (ZMMU) and the Institute of Tropical Biology Zoological Collection, Ho Chi Minh City, Vietnam (ITBCZ). Examined specimens of compared species are stored in herpetological collections of ZMMU, Zoological Institute R.A.S., St. Petersburg, Russia (ZISP), Natural History Museum,




Figure 1. Distribution of *Ophryophryne* species in southern Indochina (Langbian Plateau in Vietnam, and adjacent regions of Cambodia), indicating localities sampled in this study. Black dot in the center of an icon indicates the type locality of the new species. Locality information. **1** O Chung Chry stream, Samling Logging Concession, O'Rang Dist., Mondolkiri Prov., Cambodia (500 m a.s.l.) (Stuart et al. 2006; type locality of *O. synoria*; Mahony et al. 2017) **2** Bu Gia Map N.P., Binh Phuoc Prov., Vietnam (400 m a.s.l.) (Vassilieva et al. 2016; this study) **3** Nam Cat Tien sector of Dong Nai Biosphere Reserve, Dong Nai Prov., Vietnam (200 m a.s.l.) (Vassilieva et al. 2016; this study) **4** Loc Bac forest, Bao Loc Dist., Lam Dong Prov., Vietnam (830 m a.s.l.) (this study) **5** Cam Ly River (Ohler 2003: type locality of *O. gerti*) and Nui Ba Mt. in environs of Dalat city, Lam Dong Prov., Vietnam (ca. 1000–1800 m a.s.l.) (this study) **6** Environs of Bidoup Mt. (2000 m a.s.l.), and Giang Ly Ranger Station (1500 m a.s.l.), Bidoup–Nui Ba N.P., Lam Dong Prov., Vietnam (Poyarkov [Paiarkov] and Vassilieva 2011; this study) **7** Hon Giao Mt., Bidoup–Nui Ba N.P., Lam Dong and Khanh Hoa provincial border, Vietnam (1900–2000 m a.s.l.) (this study) **8** Hon Ba Mt., Hon Ba N.R., Dien Khanh Dist., Khanh Hoa Prov., Vietnam (950–1510 m a.s.l.) (Nguyen et al. 2014; this study) **9** Nui Chua Mt., Nui Chua N.P., Ninh Hai Dist., Ninh Thuan Prov., Vietnam (780 m a.s.l.) (this study) **10** Chu Pan Fan Mt., Chu Yang Sin N.P., Lak Dist., Dak Lak Prov., Vietnam (1900 m a.s.l.) (type locality of *Ophryophryne elfina* sp. n.; this study) **11** Chu Yang Sin Mt. environs, Krong Kmar Commune, Krong Bong Dist., Dak Lak Prov., Vietnam (700–2000 m a.s.l.) (Orlov et al. 2008; this study) **12** Tay Hoa Dist., Phu Yen Prov., Vietnam (700 m a.s.l.) (this study).

London, United Kingdom (NHMUK, formerly BMNH, though the latter acronym is retained for specimen numbers for comparability with older literature), Field Museum of Natural History, Chicago, USA (FMNH), and Yunnan University, Faculty of Biology, Kunming, China (YU).

Morphology. Specimens were photographed in life, and tissue samples for genetic analyses were taken prior to preservation, and stored in 96% ethanol. We recorded morphological data from specimens fixed and stored in 75% ethanol.

Measurements to the nearest 0.1 mm were taken using either a digital caliper, or a dissecting microscope; morphometrics of adult frogs and character terminology follows Mahony (2011a) and Mahony et al. (2013). Morphometric abbreviations are as follows:

SVL	snout to vent length;
HW	head width;
HL	head length;
ED	eye diameter;
TYD	tympanum diameter;
TYE	eye to tympanum distance;
SL	snout length;
EN	eye to narial distance;
SN	narial to snout distance;
IUE	interorbital distance, between upper eyelids;
IN	internarial distance;
UEW	upper eyelid width;
FAL	forearm length;
HAL	hand length;
FIL	first finger length;
FIIL	second finger length;
FIIL	third finger length;
FIVL	fourth finger length;
SHL	shank length;
TL	thigh length;
FOL	foot length;
TFOL	tibiotarsal articulation to tip of fourth toe distance;
IMT	inner metatarsal tubercle length.

Additionally, for the description of the type series we measured the distance between anterior orbital borders (**IFE**); distance between posterior orbital borders (**IBE**); first toe length (**TIL**); second toe length (**TIIL**); third toe length (**TIIL**); fourth toe length (**TIIVL**); fifth toe length (**TVL**). All measurements were taken on the right side of the specimen, except when a character was damaged, in which case the measurement was taken on the left side. Entire skin surface of all specimens were examined by microscope for the presence of dermal microstructures. Sex was determined by direct observation of calling in life and/or gonadal inspection by dissection.

Morphological description of larval stages included the following 15 measurements: total length (TOL); body length (BL); tail length (TAL); maximum body width (BW); maximum body height (BH); maximum tail height (TH); snout to vent length (SVL); snout to spiracle distance (SSp); maximum upper tail fin height (UF); maximum lower tail fin height (LF); internarial distance (IN); interpupilar distance (IP); rostro-narial distance (RN); nario-pupilar distance (NP); eye diameter (ED). The oral disk width and the labial tooth row formula were not recorded since in *Ophryophryne* the oral disk is modified to an extensive funnel which is closed when fixed in preservative, and oral disk structures typical for most other anurans are absent. Tadpoles were staged after Gosner (1960); morphometrics followed Grosjean (2001, 2003) and Poyarkov et al. (2015b).

All statistical analyses were performed with Statistica 6.0 (StatSoft, Inc. 2001). Morphometric characters were used for univariate analyses, corrected by body size. Sexes were separated for subsequent comparisons among the samples. One-way ANOVA and Duncan's post hoc test were used for morphometric comparisons. Multivariate statistical analyses were conducted for examination of overall morphological variation among studied populations. If some characters showed high correlation between each other, all but one of them were omitted in order to exclude the overweighting effect of these characters on the analyses. After metric values were log e-transformed, a principal component analysis (PCA) was conducted. The additional specimens of the undescribed *Ophryophryne* species, measured by LTN, were not included in the PCA to avoid potential error due to inter-observer variation of measurement techniques. A significance level of 95% was used in all statistical tests.

Comparative morphological data were obtained from museum specimens of *Ophryophryne* and (when available) photographs of these specimens in life (see Appendix 1). Data on morphology and taxonomy of *Ophryophryne* are also available from the following literature: *O. microstoma* (Bourret 1942, Liu and Hu 1962, Yang 1991, Ye et al. 1993, Fei et al. 1999, 2009, 2010, 2012, Zhang and Wen 2000, Ohler 2003, Bain et al. 2007, Yang 2008), *O. pachyproctus* (Kou 1985, Yang 1991, Ye et al. 1993, Fei et al. 1999, 2009, 2010, 2012), *O. poilani* (Bourret 1937, 1942, Ohler 2003). However, due to the considerable undiagnosed diversity within Megophryidae (Chen et al. 2017, Mahony et al. 2017), where available, we relied on the examination of type specimens, topotypic material and/or original species descriptions. Only characters verified on all specimens in the type series and referred specimens are used to represent the new species in the comparison and diagnosis sections. Specimens of *O. cf. poilani* listed in Appendix 1 were not used in the comparison of the undescribed species with *O. poilani*.

DNA isolation and sequencing. For molecular analysis, total genomic DNA was extracted from ethanol-preserved muscle or liver tissues using either standard phenol-chloroform extraction procedures (Hillis et al. 1996) followed with isopropanol precipitation (at Moscow State University: hereafter MSU), or a Qiagen DNeasy® Blood & Tissue Kit primarily following manufacturers' instructions, with the exception of an extended (10 minute) soaking step prior to the elution of extracted DNA from the column, and additional final elution step using 40 µl H₂O (at University College Dub-

lin: hereafter UCD). The isolated total genomic DNA was visualized using agarose gel electrophoresis in the presence of ethidium bromide (MSU), or SafeView™ (Applied Biological Materials Inc. — at UCD). The concentration of total DNA was measured using NanoDrop 2000 (MSU) or NanoDrop 1000 (Thermo Scientific) (UCD), and consequently either adjusted to ca. 100 ng DNA/μl (MSU), or 10 ng DNA/μl (UCD).

We amplified sequences of a continuous fragment including partial sequences of 12S rRNA and 16S rRNA genes and complete t-val gene sequence, to obtain a fragment of up to 2077 bp (base pairs) of mtDNA. For some adult specimens and larvae a partial ca. 460–500 bp fragment of the 16S rRNA gene was sequenced for molecular identification purposes. 16S rRNA is a molecular marker widely applied for biodiversity surveys in amphibians (Vences et al. 2005a, 2005b, Vieites et al. 2009), and has proven to be particularly useful in studies of megophryid diversity (Matsui et al. 2010, 2014, Ohler et al. 2011, Stuart et al. 2011, 2012, Hamidy et al. 2012, Rowley et al. 2010a, 2011a, 2012, Jiang et al. 2013, Poyarkov et al. 2015a and references therein). Amplification was performed in 25 μl reactions using either ca. 50 ng genomic DNA, 10 nmol of each primer, 15 nmol of each dNTP, 50 nmol additional MgCl₂, Taq PCR buffer (10 mM Tris-HCl, pH 8.3, 50 mM KCl, 1.1 mM MgCl₂ and 0.01% gelatine) and 1 U of Taq DNA polymerase (MSU), or 2.0 μl of genomic DNA (10 ng/μl), 2.5 μl Sigma 10x PCR buffer (excluding MgCl₂), 0.5 μl MgCl₂, 0.5 μl dNTP mix, 0.5 μl forward and reverse primer (10 ng/μl), 0.2 μl Platinum® Taq DNA Polymerase (Invitrogen) and 18.3 μl PCR grade H₂O (UCD). Primers used in PCR and sequencing were as follows: forward primers: 12SAL (AAACTGGGATTAGATACCCCACTAT; Zhang et al. 2008), LX12SN1 (TACACACCGCCCGTCA; Zhang et al. 2008), 12SA (CTGGGATTAGATACCCCACTA; Palumbi 1996), L1879 (CGTACCTTTTG-CATCATGGTC; Matsui et al. 2010), L2188 (AAAGTGGGCCTAAAAGCAGCCA; Matsui et al. 2006), 16L-1 (CTGACCGTGCAAAGGTAGCGTAATCACT; Hedges 1994); reverse primers: 16S2000H (GTGATTAYGCTACCTTTGCACGGT; Zhang et al. 2008), LX16S1R (GACCTGGATTACTCCGGTCTGAACTC; Zhang et al. 2008), 16SBr (CCGGTCTGAACT-CAGATCACGT; Palumbi 1996), H1923 (AAGTAGCTCGCTTAGTTTCGG; Matsui et al. 2010), H2317 (TTCTTGTTAC-TAGTTCTAGCAT; Shimada et al. 2011), Will6 (CCCTCGTGATGCCGTTGATAC; Wilkinson et al. 2002). Tadpoles were assigned to species based on short 16S rRNA sequences obtained using the primer pair 16L-1 (see above) and 16H-1 (CTC-CGGTCTGAACTCAGATCACGTAGG; Hedges 1994). Two Touch-Down (TD) PCR reaction protocols (Murphy and O'Brien 2007) were used: TD 63–57 for 12SA and 16SBr primers and TD 55 for all other primer pairs. Slight differences in reaction protocol were used between MSU, and UCD reactions (in parentheses). TD 55 included an initial denaturation step of 5 (2) min. at 94°C and followed with 10 cycles of denaturation for 30 (45) sec. at 96°C, primer annealing for 30 (40) sec. with annealing temperature decreasing by 1°C per cycle from 65°C to 55°C and extension step for 1 min. at 72°C, followed with 35 cycles of 30 (45) sec. at 96°C, 30 (40) sec. at 55°C and 4 (1) min. at 60°C (72°C), with the final extension step for 10 min. at 72°C. TD 63–57 consisted of 2 min. at 95°C, 6 cycles of 45 sec. at 95°C, 40 sec. at 63°C with a

reduction of 1°C each cycle, 1 min. at 72°C, followed by 35 cycles of 45 sec. at 95°C, 40 sec. at 57°C and 1 min at 72°C, and a final step of 10 min. at 72°C. PCR products were loaded onto 1% agarose gels, stained with either GelStar gel stain (Cambrex: at MSU) or SafeView™ (at UCD), and visualized in a Dark reader transilluminator (Clare Chemical). If distinct bands were produced, PCR products were purified either using 2 µl, from a 1:4 dilution of ExoSapIt (Amersham), per 5 µl of PCR product prior to cycle sequencing (MSU), or using TSAP (Promega) following manufacturers' instructions (UCD). At MSU, a 10 µl sequencing reaction included 2 µl of template, 2.5 µl of sequencing buffer, 0.8 µl of 10 pmol primer, 0.4 µl of BigDye Terminator version 3.1 Sequencing Standard (Applied Biosystems) and 4.2 µl of water. The cycle-sequencing reaction was 35 cycles of 10 sec. at 96°C, 10 sec. at 50°C and 4 min. at 60°C. Cycle sequencing products were purified by ethanol precipitation. Sequence data collection and visualization were performed on an ABI 3730xl automated sequencer (Applied Biosystems). At UCD, purified PCR products were Sanger sequenced in both directions by Macrogen (Europe). The forward and reverse sequences were checked visually either in Chromas Pro software (Technelysium Pty Ltd., Tewantin, Australia: at MSU) and a consensus sequence was compiled with BioEdit 5.0.9 (Hall 1999: at MSU), or using CodonCodeAligner 3.7.1 (CodonCode Corporation, Dedham, Massachusetts: at UCD). Sequences were submitted to a BLAST search in GenBank to confirm that the intended sequences had been amplified. The obtained sequences are deposited in GenBank under the accession numbers KY425352–KY425411 and KY515232–KY515233 (see Table 1).

Phylogenetic analyses. Sequences coding for the 12S rRNA–16S rRNA mtDNA genes of 66 megophryid specimens: 53 *Ophryophryne*, representing all currently recognized species, and outgroup sequences of two *Brachytarsophrys* species, eight *Megophrys s. lato* species (including seven *Xenophrys* and one *Megophrys s. stricto* species), two *Leptobrachium* Tschudi, 1838, and one *Leptolalax* Dubois, 1980 species (Table 1), were included in the final alignment and subjected to phylogenetic analyses. Nucleotide sequences were initially aligned using ClustalX 1.81 (Thompson et al. 1997) with default parameters, and then optimized manually in BioEdit 7.0.5.2 (Hall 1999) and MEGA 6.0 (Tamura et al. 2013). Mean uncorrected genetic distances (*p*-distances) between sequences were determined with MEGA 6.0 (Tamura et al. 2013); the existence of “barcode gap” was estimated using the online version of ABGD (Puillandre et al. 2012). MODELTEST v.3.06 (Posada and Crandall 1998) was used to estimate the optimal evolutionary models to be used for the data set analysis. The best-fitting model as suggested by the Akaike Information Criterion (AIC) was the general time-reversible (GTR) model of DNA evolution with a gamma shape parameter (G).

Maximum Likelihood (ML) analysis was conducted using Treefinder (Jobb et al. 2004). Transitions and transversions were equally weighted, and gaps were treated as missing data. Confidence in tree topology was tested by non-parametric bootstrap analysis (Felsenstein 1985) with 1000 replicates. Bayesian inference (BI) was conducted using MrBayes 3.1.2 (Huelsenbeck and Ronquist 2001, Ronquist and Huelsenbeck 2003); Metropolis-coupled Markov chain Monte Carlo (MCMCMC) analyses were

Table 1. Specimens and GenBank sequences of *Ophryophryne* and outgroup Megophryidae representatives used in molecular analyses. AN – Accession number. Numbers of localities (No. 1–12) correspond to those in Figures 1 and 2. For detailed specimen information see Appendix 1. Asterisk marks holotype specimen, double asterisk marks topotype specimens (continues on next two pages). Museum abbreviations for the specimens from which sequences were generated in this study are explained in the Materials and methods section.

GenBank AN	Voucher ID	Species	Locality	Elevation (m a.s.l.)	Reference
KY022198	FMNH 262778*	<i>O. synoria</i>	1 – Cambodia, Mondolkiri Prov., O'Reang	500	Mahony et al. 2017
KY425353	ZMMU ABV-00379	<i>O. synoria</i>	2 – Vietnam, Binh Phuoc Prov., Bu Gia Map N.P., Dac Ca River	400	this paper
KY425354	ZMMU NAP-00731	<i>O. synoria</i>	2 – Vietnam, Binh Phuoc Prov., Bu Gia Map N.P., Dac Ca River	400	this paper
KY425355	ZMMU ABV-00380	<i>O. synoria</i>	2 – Vietnam, Binh Phuoc Prov., Bu Gia Map N.P., Dac Ca River	400	this paper
KY425356	ZMMU ABV-00376	<i>O. synoria</i>	2 – Vietnam, Binh Phuoc Prov., Bu Gia Map N.P., Dac Ca River	400	this paper
KY425357	ZMMU NAP-00834	<i>O. synoria</i>	3 – Vietnam, Dong Nai Prov., Nam Cat Tien N.P., Da Ta Po River	200	this paper
KY425358	ZMMU ABV-00209	<i>O. synoria</i>	4 – Vietnam, Lam Dong Prov., Bao Loc, Loc Bac forestry	830	this paper
KY425359	ZMMU ABV-00159	<i>O. synoria</i>	4 – Vietnam, Lam Dong Prov., Bao Loc, Loc Bac forestry	830	this paper
KY425360	ZMMU NAP-01756	<i>O. synoria</i>	6 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Giang Ly	1500	this paper
KY425361	ZMMU NAP-01787	<i>O. synoria</i>	6 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Giang Ly	1500	this paper
KY425362	ZMMU NAP-01835	<i>O. synoria</i>	6 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Giang Ly	1500	this paper
KY425363	ZISP NLO-36349	<i>O. synoria</i>	11 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P., Chu Yang Sin Mt.	1000	this paper
KY425364	ZISP NLO-36554	<i>O. synoria</i>	11 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P., Chu Yang Sin Mt.	1000	this paper
KY425365	KIZ-013663**	<i>O. gerti</i>	5 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Langbian Mt.	1800	this paper
KY425366	KIZ-013664**	<i>O. gerti</i>	5 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Langbian Mt.	1800	this paper
KY425367	KIZ-013662**	<i>O. gerti</i>	5 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Langbian Mt.	1800	this paper
KY425368	ZMMU NAP-01878	<i>O. gerti</i>	6 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Giang Ly	1500	this paper
KY425369	ZMMU NAP-01789	<i>O. gerti</i>	6 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Giang Ly	1500	this paper
KY425370	ZMMU NAP-02471	<i>O. gerti</i>	6 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Giang Ly	1500	this paper
KY425371	ZMMU NAP-01790	<i>O. gerti</i>	6 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Giang Ly	1500	this paper
KY425372	ZMMU NAP-01788	<i>O. gerti</i>	6 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Giang Ly	1500	this paper
KY425373	ZMMU NAP-02758	<i>O. gerti</i>	11 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P.	1000	this paper
KY425374	ZMMU ABV-00530	<i>O. gerti</i>	11 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P.	1000	this paper
KY425375	ZMMU NAP-02759	<i>O. gerti</i>	11 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P.	1000	this paper

GenBank AN	Voucher ID	Species	Locality	Elevation (m a.s.l.)	Reference
KY425376	ZMMU ABV-00577	<i>O. gerti</i>	11 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P.	1000	this paper
KY425377	ZMMU NAP-02760	<i>O. gerti</i>	11 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P.	1000	this paper
KY425378	ZISP NLO-36510	<i>O. gerti</i>	11 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P.	1000	this paper
KY425379	ZMMU ABV-00454	<i>O. elfina</i> sp. n.	6 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P.; Bidoup Mt.	2000	this paper
KY425380	ZMMU ABV-00455	<i>O. elfina</i> sp. n.	6 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P.; Bidoup Mt.	2000	this paper
KY515233	ZMMU NAP-01169	<i>O. elfina</i> sp. n. (larva)	6 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P.; Bidoup Mt.	2000	this paper
KY425381	ZMMU NAP-01782	<i>O. elfina</i> sp. n.	7 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P.; Hon Giao Mt.	2000	this paper
KY425382	ZMMU NAP-01783	<i>O. elfina</i> sp. n.	7 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P.; Hon Giao Mt.	2000	this paper
KY425383	ZMMU NAP-01757	<i>O. elfina</i> sp. n.	7 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P.; Hon Giao Mt.	2000	this paper
KY425384	ZMMU NAP-01758	<i>O. elfina</i> sp. n.	7 – Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P.; Hon Giao Mt.	2000	this paper
KY425385	ZMMU ABV-00316	<i>O. elfina</i> sp. n.	8 – Vietnam, Khanh Hoa Prov., Hon Ba N.R., Hon Ba Mt.	1500	this paper
KY425386	KIZ YPX-05429	<i>O. elfina</i> sp. n.	9 – Vietnam, Ninh Thuan Prov., Nui Chua N.P.	780	this paper
KY425387	KIZ YPX-05457	<i>O. elfina</i> sp. n.	9 – Vietnam, Ninh Thuan Prov., Nui Chua N.P.	780	this paper
KY425388	KIZ YPX-05428	<i>O. elfina</i> sp. n.	9 – Vietnam, Ninh Thuan Prov., Nui Chua N.P.	780	this paper
KY425389	ZMMU NAP-02658*	<i>O. elfina</i> sp. n.	10 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; Chu Pan Fan Mt.	1900	this paper
KY515232	ZMMU NAP-02673**	<i>O. elfina</i> sp. n. (larva)	10 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; Chu Pan Fan Mt.	1900	this paper
KY425390	ZISP NLO-36522	<i>O. elfina</i> sp. n.	11 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; Chu Yang Sin Mt.	2000	this paper
KY425391	ZMMU ABV-00581	<i>O. elfina</i> sp. n.	11 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; Chu Yang Sin Mt.	1800	this paper
KY425392	ZMMU ABV-00580	<i>O. elfina</i> sp. n.	11 – Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; Chu Yang Sin Mt.	1975	this paper
KY425393	DVT-00393	<i>O. elfina</i> sp. n.	12 – Vietnam, Phu Yen Prov., Tay Hoa	700	this paper
KY022203	AMNH A163680	<i>O. hansi</i>	Vietnam, Quang Nam Prov., Tra My, Tra Don	930	Mahony et al. 2017
DQ283377	AMNH A163669	<i>O. hansi</i>	Vietnam, Quang Nam Prov., Tra My, Tra Don	970	Frost et al. 2006
KY425395	ZMMU NAP-06485	<i>O. hansi</i>	Vietnam, Kon Tum Prov., Thac Nham forest	1100	this paper
KY425396	ZMMU NAP-06501	<i>O. hansi</i>	Vietnam, Gia Lai Prov., Kon Chu Rang N.R.	1000	this paper
KY425397	ZMMU NAP-06524	<i>O. hansi</i>	Vietnam, Gia Lai Prov., Kon Chu Rang N.R.	1000	this paper
KY425398	ZMMU NAP-06502	<i>O. hansi</i>	Vietnam, Gia Lai Prov., Kon Chu Rang N.R.	1000	this paper
KY425399	ZMMU NAP-06525	<i>O. hansi</i>	Vietnam, Gia Lai Prov., Kon Chu Rang N.R.	1000	this paper
KY022200	KUH 311601	<i>O. microstoma</i>	China, Guangxi Prov., Shiwan Dashed N.R., Fulong	500	Mahony et al. 2017
KY022199	AMNH A168682	<i>O. microstoma</i>	Vietnam, Lao Cai Prov., Van Ban Dist., Nam Tha	330	Mahony et al. 2017

GenBank AN	Voucher ID	Species	Locality	Elevation (m a.s.l.)	Reference
KY022201	AMNH A163668	<i>O. cf. poilani</i>	Vietnam, Quang Nam Prov., Tra My, Tra Don	980	Mahony et al. 2017
KY022202	AMNH A169287	<i>O. cf. poilani</i>	Vietnam, Thua Thien–Hue Prov., A Luoi Dist., A Roang	680	Mahony et al. 2017
JX564854	ZP-AM 44	<i>Brachytarsophrys carinense</i>	–	–	Zhang et al. 2013
KY425404	ZMMU NAP-06324	<i>Brachytarsophrys feae</i>	Vietnam, Vinh Phuc Prov., Tam Dao	–	this paper
KY425405	ZMMU NAP-03994	<i>Xenophrys cf. acutus</i>	Thailand, Satun Prov.	–	this paper
KY425406	ZMMU NAP-05005	<i>Xenophrys cf. dauvinmontis</i>	Vietnam, Dien Bien Prov., Muong Nhe, Muong Nhe N.R.	–	this paper
KY425407	ZMMU NAP-04137	<i>Xenophrys cf. parva</i>	Thailand, Surathani Prov.	–	this paper
KY425408	ZMMU NAP-04423	<i>Xenophrys cf. maosomensis</i>	Vietnam, Dien Bien Prov., Muong Nhe, Muong Nhe N.R.	–	this paper
AY561308	ZYCI 500	<i>Xenophrys minor</i>	China	–	Zheng et al. 2004
AY561307	ZYCI 513	<i>Xenophrys omeimontis</i>	China, Sichuan	–	Zheng et al. 2004
KY425409	DVT-04135	<i>Xenophrys</i> sp.	Vietnam, Lao Cai Prov.	–	this paper
KY425410	ZMMU NAP-05095	<i>Megophrys nasuta</i>	Malaysia, Sarawak	–	this paper
KJ630505	SCUM120630	<i>Leptobrachium boringii</i>	China, Sichuan Prov., Emei Shan Mt.	–	Xu et al. 2014
KY425411	DVT-00298	<i>Leptobrachium banuae</i>	Vietnam, Phu Yen Prov., Tay Hoa	–	this paper
JX564874	MVZ-Herp-223642	<i>Leptoldax cf. pelodytoides</i>	–	–	Zhang et al. 2013

run with one cold chain and three heated chains for four million generations and sampled every 1,000 generations. Five independent MCMCMC runs were performed and 1,000 trees were discarded as burn-in. Confidence in tree topology was assessed by posterior probability (Huelsenbeck and Ronquist 2001).

We *a priori* regarded tree nodes with bootstrap (BS) values 75% or greater and posterior probabilities (BPP) values over 0.95 as sufficiently resolved, BS values between 75% and 50% (BPP between 0.95 and 0.90) were regarded as tendencies, and BS values below 50% (BPP below 0.90) were considered to be unresolved (Huelsenbeck and Hillis 1993).

Acoustic analyses. Calls were recorded using a portable digital audio recorder Zoom h4n (ZOOM Corporation, Tokyo, Japan) in stereo mode with 96 kHz sampling frequency and 16-bit precision, or using a Marantz 660 digital tape recorder (D&M Professional, Kanagawa, Japan) in mono mode with sampling rate at 48 kHz and 16-bit precision with a high-sensitivity Sennheiser K6 ME66 cardioid electret condenser microphone (Sennheiser electronic, Wedemark, Germany), or using a Nikon D 600 digital SLR camera (Nikon Corporation, Japan) in video mode with audio tracks removed from video recordings using Avisoft SASLab Pro software v. 5.2.05 (Avisoft Bioacoustics, Germany) with a 48 kHz sampling frequency and 16-bit precision. Temperature was measured at the calling sites immediately after audio recording using a digital thermometer, KTJ TA218A Digital LCD Thermometer-Hydrometer. All recordings were made in situ in the natural habitats of respective specimens. Advertisement calls of the undescribed *Ophryophryne* species were recorded on the mountain summit of Bidoup in the Bidoup-Nui Ba National Park (hereafter N.P.), Lam Dong Prov., eastern edge of the Langbian Plateau, Vietnam (12°06'42.4"N; 108°39'33.6"E, 1930–1940 m a.s.l.), on 10 and 15 April 2014, and 10 February 2015, between 16:05–18:35 h and at temperatures from 11.3°C in February to 17.5°C in April. In total, we made five recordings from three vocalizing males. Advertisement calls of *O. gerti* were recorded in Chu Yang Sin N.P., Dak Lak Prov., northern edge of the Langbian Plateau, Vietnam (12°24'01.6"N; 108°21'11.0"E, 1020 m a.s.l. and 12°25'25.7"N; 108°21'52.5"E, 1040–1045 m a.s.l.), on 22–27 May 2014, between 20:40–23:10 h at 22–22.5°C. In total, we made three recordings from three males. Advertisement calls of *O. synoria* were recorded in Chu Yang Sin N.P., Dak Lak Prov., Tay Nguyen region, Vietnam (12°28'0.94"N; 108°20'45.4"E, 700–800 m a.s.l.) on 25 May 2008, between 21:56–22:30 h at 21°C. In total, three recordings from three males were made.

All recordings were standardized by Avisoft SASLab Pro software v. 5.2.05 in mono format with sampling rate at 48 kHz and 16-bit precision, and low-frequency noise was reduced using the low-pass filter (up to 1000 Hz). Calls were analyzed using Avisoft SASLab Pro software v. 5.2.05; all parameters were measured using the reticule and standard cursors in the spectrogram window of Avisoft. Spectrograms for analyses were created using the Hamming window, FFT-length 1024 points, frame 100%, and overlap 87.5%. Figure spectrograms were created using the Hamming window, FFT-length 512 points, frame 100%, and overlap 75%. In total, we measured 1797 calls of the new *Ophryophryne* sp., 533 calls of *O. gerti* and 200 calls of *O. synoria*.

Four temporal parameters were measured: the duration of each call, the interval between successive calls within each series, the duration of series, the interval between successive series, and five frequency parameters: the initial and final fundamental frequency, the minimum and maximum of fundamental frequency and the frequency of maximum amplitude (also “F peak”). Then we calculated the frequency range as the difference between the maximum and minimum of fundamental frequencies and the call repetition rate per recording/series (calls/s) for each recording/series as a ratio of number of all calls within the recording/series (excluding series consisting of just one call) to recording/series duration. All numerical parameters are given as mean \pm SE, the minimum and maximum values are given in parentheses (min–max).

To compare acoustic characteristics between three species of *Ophryophryne* we applied one-way ANOVA with Tukey post hoc for the values of the parameters for which distributions did not differ from normality ($p > 0.05$, Kolmogorov–Smirnov test). Otherwise we used nonparametric Kruskal–Wallis ANOVA with Mann–Whitney U post hoc test.

The records of advertisement calls were deposited at the Fonoteca Zoologica and are available at the website <http://www.fonozoo.com> (under the accession numbers 9954–9964).

Results


Sequence data

The final alignment of the studied 12S rRNA–16S rRNA mtDNA gene fragment consisted of 2077 sites: 1439 sites were conserved and 567 sites were variable, of which 465 were found to be parsimony-informative. The transition–transversion bias (R) was estimated as 2.06. Nucleotide frequencies were A = 32.8%, T = 27.6%, C = 21.6%, and G = 17.9% (all data given for ingroup only).

Phylogenetic relationships and geographic distribution of mtDNA haplotypes

We achieved high resolution of phylogenetic relationships among taxa within *Ophryophryne*, with all major nodes fully resolved (BPP = 1.0; BS = 100%; Fig. 2). Monophyly of species-level groups and species complexes in *Ophryophryne* were also significantly supported (BS > 90%; BPP \geq 0.95). However, phylogenetic relationships between the taxa of outgroup Megophryinae are poorly resolved with major nodes in the tree having low or insignificant levels of support (BPP < 0.95; BS < 75%). Bayesian and Maximum Likelihood analyses resulted in essentially similar topologies (see Fig. 2) slightly differing from each other only in associations for several poorly supported outgroup nodes.

Our analyses (Fig. 2) inferred the following set of phylogenetic relationships among studied megophryid taxa:



Our data confirm the monophyly of Megophryinae with respect to outgroup taxa (*Leptobrachium* and *Leptolalax*) (1.0/100; hereafter node support values are given for BPP/BS respectively). Within Megophryinae, the sample of *Megophrys nasuta* (Schlegel, 1858), representing the genus *Megophrys* s. *stricto*, forms the most basal split; this lineage is recovered as a sister group with respect to all other Megophryinae. Phylogenetic relationships among other genera of Megophryinae remain essentially unresolved; while monophyly of *Brachytarsophrys* received high support (0.99/100), species assigned to *Xenophrys* form two weakly supported groups, paraphyletic with respect to *Ophryophryne* and *Brachytarsophrys*.

Monophyly of the genus *Ophryophryne* is strongly supported by all analyses (1.0/100). General topology of the Bayesian tree suggests that the genus *Ophryophryne* is divided into two major groups: the first group joins taxa from southern China, northern and central Indochina (Group I, see Fig. 2), while the second group comprises lineages of *Ophryophryne* confined to the Langbian Plateau in southern Vietnam and adjacent Cambodia (Group II, see Fig. 2). Both clades are reciprocally monophyletic with high support values (1.0/100).

Within Group I, the clade consisting of two specimens from central Vietnam (AMNH A-169287, Thua Thien-Hue Prov., and AMNH A-163668, Quang Nam Prov.: identified as *O. gerti* by Bain et al. 2007) forms a sister clade to all other species in this group (1.0/100). This lineage is determined to be distantly related to the topotype specimens of *O. gerti* from the Langbian Plateau, and based on examination of these specimens we tentatively regard these specimens as *O. cf. poilani* (see discussion below).

Within Group I, specimens of *O. microstoma* from southern China (KUH 311601) and northern Vietnam (AMCC 141231) are clustered together (1.0/100) forming a sister clade to a group comprising specimens identified as *O. hansii* Ohler, 2003, from Kon Tum Plateau in central Vietnam; monophyly of the latter group is also strongly supported (1.0/100) (Fig. 2).

Within Group II, Subclade A joins the medium-sized specimens from environs of the type locality of *O. gerti* (Langbian Mt., environs of Da Lat city, Lam Dong Prov., Vietnam, up to 1800 m a.s.l.; see Fig. 1: Loc. 5) with two other populations found at medium elevation (1000–1500 m a.s.l.) on the eastern (Bidoup–Nui Ba N.P., Lam Dong Prov., Vietnam, see Fig. 1: Loc. 6) and northern (Chu Yang Sin N.P., Dak Lak Prov., Vietnam; see Fig. 1: Loc. 11) edges of the Langbian Plateau (Fig. 2). In the present paper we treat this lineage as the species level clade representing *O. gerti* s. *stricto* (see discussion below).

The second species level clade comprises large-sized *Ophryophryne* from comparatively lowland populations in the western foothills of the Langbian Plateau (Fig. 1, Locs 1–4, < 1000 m a.s.l.) and large-sized *Ophryophryne* from medium elevations in the northern parts of the plateau (Fig. 1, Locs 6 and 11; 1000–1500 m a.s.l.). Two reciprocally monophyletic (1.0/100) subclades are revealed in this clade (Fig. 2). Subclade B joins montane populations from 1000–1500 m a.s.l. (Fig. 1, Locs 4, 6 and 11) and a recently discovered lowland (200 m a.s.l.) population from Cat Tien N.P. in Dong Nai Prov. Subclade C joins two populations from the westernmost edge of the Langbian Plateau, which include the holotype of *O. synoria* from Mondolkiri Prov. in eastern

Cambodia (Fig. 1, Loc. 1; 500 m a.s.l.) and a population from Bu Gia Map N.P. in the adjacent Vietnamese province of Binh Phuoc (Fig. 1, Loc. 2; 400 m a.s.l.). These localities are close to each other and geographically belong to one hilly region on the western edge of the Langbian Plateau. Based on phylogenetic and morphological data we herein regard both subclades B and C as the species level clade representing *O. synoria*.

The third species level clade forms a sister clade with respect to a clade comprised of *O. gerti s. stricto* and *O. synoria* (Fig. 2). It joins small-sized *Ophryophryne* specimens, all collected from both high elevations (> 1750 m a.s.l.) in the northern and eastern parts of the Langbian Plateau (see Fig. 1: Bidoup and Hon Giao Mts., Lam Dong Prov., Locs 6–7; and Chu Pan Fan and Chu Yang Sin Mts., Dak Lak Prov., Locs 10–11), and from lower elevations (700–1510 m a.s.l.) on the summits of three mountains representing the easternmost outcrops of the Langbian Plateau (see Fig. 1: Hon Ba Mt., Khanh Hoa Prov., Loc. 8; Nui Chua Mt., Ninh Thuan Prov., Loc. 9; Tay Hoa, Phu Yen Prov., Loc. 12). Among these populations, samples from the summit of Nui Chua Mt. (Fig. 1, Loc. 9) form Subclade D (Fig. 2), forming a sister clade with respect to all other populations (Fig. 2, Subclade E; monophyly support 0.95/98). This clade of small-sized *Ophryophryne* from the northern and eastern parts of the Langbian Plateau currently cannot be assigned to any of the recognized species and represents a new species described herein.

Intra- and interspecific differentiation of mtDNA haplotypes

The observed interspecific sequence divergence within the genus *Ophryophryne* varied from $p = 4.1\%$ to $p = 13.0\%$ (Table 2). The values of uncorrected genetic p -distances in ingroup and outgroup comparisons slightly overlapped: sequence divergence between *Ophryophryne* and outgroup taxa varied from $p = 8.8\%$ to $p = 24.7\%$. The minimal interspecific p -distance between recognized nominal species in our analysis was found between the sister species *O. gerti* and *O. synoria* ($p = 4.1\%$ – 5.0%). The maximum p -distance for *Ophryophryne* was observed between *O. synoria* and *O. hansii* ($p = 12.6\%$ – 13.0%) (see Table 2). The ABGD analysis revealed the existence of a “barcoding gap” at genetic distance value $p = 4.0\%$ in the 16S rRNA gene.

Intraspecific distances within *Ophryophryne* species in our analysis varied from $p = 0.5\%$ (in *O. gerti*), to $p = 3.3\%$ among two samples of *O. cf. poilani* from Quang Nam and Thua Thien-Hue provinces of Vietnam, and to $p = 3.7\%$ among two samples of *O. microstoma* from China and Vietnam respectively. The latter two values are higher than usual intraspecific distances in the 16S rRNA gene in Anura (Vences et al. 2005a, 2005b, Vieites et al. 2009); we recognize that identification of these lineages as conspecific is preliminary, based on morphology and topology of the mtDNA tree. Further studies are needed to clarify their taxonomic status. We also found significant genetic differentiation between intraspecific lineages in two of the three *Ophryophryne* species inhabiting the Langbian Plateau. Sequence divergence between Subclades B and C of *O. synoria* is $p = 2.6\%$, while the differentiation between the Nui Chua population (Subclade D) and all other populations of the new species (Subclade E) was even greater at $p = 3.1\%$.

The newly discovered lineage of *Ophryophryne* from highlands of the northern and eastern parts of the Langbian Plateau was found to have the lowest genetic distance with respect to *O. gerti* ($p = 8.2\%–9.1\%$). This value is much higher than the minimum genetic distances observed in intraspecific comparisons between species of *Ophryophryne* included in this study (Table 2).

Morphological differentiation

Among the three species examined, mean SVL varied significantly, ranging from 26.9 to 53.7 mm in males and from 35.1 to 70.7 mm in females (Table 3). For SVL, *post hoc* analyses of one-way ANOVA revealed that males were significantly smaller than females in all three species of Langbian *Ophryophryne* (one-way ANOVA, $p < 0.05$; Duncan test, $p < 0.05$). Body size variation among adult males and females of Langbian *Ophryophryne* is shown in Figure 3. All three species are clearly different in body size, with *O. synoria* being the largest, and the undescribed *Ophryophryne* species being the smallest species known for the genus (male SVL values overlap with values for *O. pachyproctus*, N.A. Poyarkov, pers. observ.). *Ophryophryne gerti* occupies an intermediate position between these two species, with SVL values of males (31.7–42.2 mm) slightly overlapping both with those for *O. synoria* (38.2–53.7 mm) and the undescribed *Ophryophryne* species (26.9–33.9 mm).

The results of the multivariate PCA-analysis of the morphometric data are shown in Fig. 4 (data given for males only). The discriminative power of PCA factors derived from analysis of morphometric characters is shown in Appendix 2. For males, F1 explained 77.80% of the variability and F2 explained 5.99%. The two-dimensional plots of both the first two principal components (Factor 1 and Factor 2; Fig. 4A) and the first vs. the third principal components (Factor 1 and Factor 3; Fig. 4B) for males completely discriminated the following three morpho-groups: (I) *O. gerti*, (II) *O. synoria*, and (III) the undescribed *Ophryophryne* species. Our multivariate analysis included meristic data for the holotypes of two *Ophryophryne* species: *O. synoria* (FMNH 262779, male) and *O. gerti* (BMNH 1921.4.1.324, male). Both holotypes were assigned correctly to the respective groups, representing *O. synoria* and *O. gerti*, in full concordance with the results of molecular analyses (Fig. 4).

In summary, both in body size and other meristic characters, all three examined lineages of Langbian *Ophryophryne* form clearly separated morphological groups, also supported by multivariate statistical analysis. The small-sized population of the undescribed *Ophryophryne* species appears to be morphologically distinct from sympatric *O. gerti* and *O. synoria*, as well as from other congeners (see Comparisons for details).

Acoustic differentiation

Measurements of advertisement call parameters for three *Ophryophryne* species found on the Langbian Plateau are given in Table 4, oscillograms and sonograms are given

Table 2. Uncorrected p -distance (percentage) between 16S rRNA sequences of *Ophryophryne* species and outgroup species included in phylogenetic analyses (below the diagonal), and standard error estimates (above the diagonal). The ingroup mean uncorrected p -distances are shown on the diagonal and shaded with grey.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
1 <i>O. gerti</i> (Subclade A)	0.5	1.1	1.1	1.7	1.6	1.9	1.8	1.7	2.0	1.9	1.9	2.1	2.2	2.1	2.2	2.2	2.1	1.9	2.2	2.4
2 <i>O. synoria</i> (Subclade B)	4.1	0.4	0.9	1.6	1.6	1.9	1.8	1.8	1.8	1.8	1.7	2.0	2.1	2.1	2.2	2.2	2.0	1.9	2.2	2.3
3 <i>O. synoria</i> (Subclade C)	5.0	2.6	0.9	1.7	1.6	1.8	1.9	1.7	1.8	1.8	1.6	2.0	2.1	2.1	2.1	2.1	2.0	1.9	2.2	2.2
4 <i>O. elfina</i> sp. n. (Subclade D)	9.1	9.3	10.0	0.0	0.9	1.8	1.6	1.7	2.1	1.8	1.9	2.2	2.0	2.2	2.2	2.3	2.1	2.1	2.3	2.2
5 <i>O. elfina</i> sp. n. (Subclade E)	8.2	8.9	8.7	3.1	0.8	1.8	1.6	1.6	2.1	2.0	2.1	2.2	1.8	2.3	2.2	2.1	2.1	2.0	2.3	2.3
6 <i>O. microstoma</i>	10.9	11.8	11.6	9.7	9.5	3.7	1.3	1.5	1.6	1.9	1.7	1.7	1.8	1.9	2.1	2.2	2.0	1.9	2.4	2.4
7 <i>O. hans</i>	10.9	12.6	13.0	9.7	10.0	7.6	0.0	1.3	1.7	1.8	1.7	1.9	1.8	2.1	2.2	2.3	1.9	1.8	2.4	2.5
8 <i>O. cf. poilani</i>	9.1	10.5	10.6	8.6	9.4	7.8	7.3	3.3	1.8	1.7	1.7	1.8	1.7	2.0	2.1	2.5	1.9	1.7	2.2	2.2
9 <i>Xenophrys minor</i>	12.8	13.0	12.7	10.3	10.9	8.0	10.8	8.8	—	1.8	1.4	1.9	2.0	2.0	2.2	2.1	1.5	1.7	2.3	2.4
10 <i>Xenophrys</i> sp.	11.6	12.2	11.9	11.1	11.7	10.3	9.5	8.8	7.8	—	1.3	1.8	1.8	1.8	2.2	2.4	1.7	1.7	2.2	2.2
11 <i>Xenophrys omeimontis</i>	10.8	11.4	11.0	9.9	10.2	8.8	9.6	8.4	5.3	3.7	—	1.7	1.8	1.8	2.1	2.1	1.2	1.6	2.2	2.2
12 <i>Xenophrys</i> cf. <i>maosonensis</i>	13.6	14.7	15.1	14.4	15.2	10.5	12.8	9.7	11.1	10.3	9.9	—	1.4	1.4	1.9	2.2	1.8	1.7	2.4	2.4
13 <i>Xenophrys</i> cf. <i>parva</i>	13.6	14.2	14.3	11.1	10.7	10.1	11.4	9.7	9.5	9.1	9.1	7.4	—	1.6	1.9	2.3	1.7	1.8	2.5	2.4
14 <i>Xenophrys</i> cf. <i>dauvetmontis</i>	12.8	13.4	14.3	14.8	14.5	11.3	14.0	11.5	11.1	9.5	9.1	6.6	6.2	—	2.0	2.3	2.0	2.1	2.6	2.3
15 <i>Xenophrys</i> cf. <i>aceras</i>	16.5	16.3	16.0	18.1	18.3	16.9	18.9	14.8	14.8	14.0	12.8	13.6	14.4	13.6	—	2.5	2.0	1.9	2.3	2.4
16 <i>Megophrys nasuta</i>	14.5	16.0	16.4	16.0	14.5	15.8	17.8	15.8	14.0	14.4	13.6	13.6	13.2	14.8	16.5	—	2.0	2.2	2.6	2.4
17 <i>Brachytarsophrys feae</i>	12.8	13.0	13.7	10.7	11.5	11.5	11.7	10.9	7.0	7.4	5.3	11.1	9.1	11.1	12.3	14.4	—	1.4	2.4	2.4
18 <i>Brachytarsophrys carinense</i>	11.5	11.4	12.0	11.5	11.5	11.5	11.3	10.1	8.6	7.4	7.4	10.3	9.5	11.9	12.3	12.8	4.9	—	2.2	2.3
19 <i>Lepidolax</i>	17.5	18.8	19.4	18.5	19.9	21.0	21.1	18.9	21.8	19.8	21.0	20.2	19.3	21.4	22.6	23.9	20.2	18.9	—	2.2
20 <i>Leptobrachium</i>	23.1	23.9	24.0	22.2	21.8	23.5	24.7	23.1	22.8	22.0	21.8	23.3	22.4	23.7	23.7	22.2	20.4	20.8	21.8	—

Table 3. Measurements of the three species of *Ophryophryne* found on the Langbian Plateau (southern Vietnam) and in adjacent Cambodia. For abbreviations see Material and methods. Values are given as means ± SE (min–max), *N* — number of specimens. All measurements are given in mm (continues on next page).

Species	SVL	HW	HL	ED	TYD	TYE	SL	EN	NS	IUE	IN	UEW
<i>Ophryophryne gerti</i>												
Males	35.9 ± 3.4 (31.7–42.2)	10.3 ± 0.8 (9.2–12.0)	10.2 ± 0.9 (9.0–12.0)	4.4 ± 0.4 (3.8–5.3)	2.4 ± 0.4 (1.8–3.1)	2.0 ± 0.3 (1.7–2.5)	3.2 ± 0.5 (2.6–3.8)	1.4 ± 0.3 (0.7–1.8)	1.6 ± 0.3 (1.1–2.2)	3.3 ± 0.5 (2.2–4.1)	2.7 ± 0.4 (1.9–3.4)	3.1 ± 0.4 (2.2–3.8)
Females	45.1 ± 2.2 (43.1–47.4)	11.3 ± 0.4 (10.9–11.7)	11.4 ± 0.6 (10.9–12.1)	4.8 ± 0.4 (4.4–5.1)	2.7 ± 0.2 (2.5–3.0)	2.5 ± 0.2 (2.3–2.6)	3.4 ± 0.4 (2.9–3.6)	1.6 ± 0.3 (1.2–1.9)	1.7 ± 0.4 (1.4–2.2)	3.8 ± 0.2 (3.6–4.0)	2.8 ± 0.1 (2.7–2.9)	3.9 ± 0.3 (3.5–4.2)
<i>Ophryophryne elfina</i> sp. n.												
Males	29.7 ± 1.8 (26.9–33.9)	8.6 ± 0.7 (7.2–10.1)	8.9 ± 0.6 (7.4–9.8)	3.7 ± 0.3 (3.1–4.3)	2.2 ± 0.2 (1.6–2.7)	1.5 ± 0.2 (1.1–2.1)	2.7 ± 0.4 (2.0–3.3)	1.2 ± 0.2 (0.8–1.6)	1.4 ± 0.3 (1.0–2.2)	2.7 ± 0.3 (2.1–3.4)	2.3 ± 0.5 (1.5–3.3)	2.9 ± 1.0 (2.4–7.9)
Females	35.6 ± 0.5 (35.1–36.5)	9.7 ± 0.5 (9.0–10.2)	9.8 ± 1.1 (8.6–11.0)	4.0 ± 0.2 (3.6–4.2)	2.3 ± 0.4 (1.9–3.0)	1.7 ± 0.1 (1.6–1.9)	3.1 ± 0.5 (2.4–3.6)	1.4 ± 0.3 (1.0–1.7)	1.7 ± 0.3 (1.2–2.1)	2.9 ± 0.5 (2.3–3.6)	2.5 ± 0.4 (1.7–2.9)	2.9 ± 0.2 (2.6–3.1)
<i>Ophryophryne synoria</i>												
Males	45.7 ± 4.3 (38.2–53.7)	13.8 ± 0.9 (12.5–15.6)	13.5 ± 1.0 (12.2–15.4)	5.1 ± 0.3 (4.7–5.6)	3.5 ± 0.5 (3.0–4.7)	2.8 ± 0.4 (1.9–3.6)	3.8 ± 0.5 (3.0–4.6)	1.9 ± 0.4 (1.3–2.5)	1.6 ± 0.3 (1.2–2.2)	4.5 ± 0.6 (3.5–5.5)	3.1 ± 0.3 (2.5–3.5)	3.7 ± 0.4 (3.0–4.1)
Females	60.8 ± 9.7 (51.4–70.7)	16.6 ± 1.8 (14.6–18.0)	16.1 ± 1.8 (14.0–17.4)	5.3 ± 0.4 (5.0–5.7)	3.7 ± 0.2 (3.4–3.9)	3.3 ± 0.3 (3.0–3.6)	4.0 ± 0.3 (3.7–4.3)	2.0 ± 0.4 (1.6–2.4)	1.8 ± 0.4 (1.3–2.2)	4.7 ± 0.2 (4.4–4.9)	3.5 ± 0.6 (3.0–4.1)	4.1 ± 0.2 (3.8–4.2)

Table 3 Continued. Measurements of the three species of *Ophryophryne* found on the Langbian Plateau (southern Vietnam) and in adjacent Cambodia. For abbreviations see Material and methods. Values are given as means \pm SE (min–max), *N* — number of specimens. All measurements are given in mm.

Species	EAL	HAL	FIL	FIIL	FIHL	FIVL	SHL	TL	FOL	TFOL	IMT
<i>Ophryophryne gerti</i>											
Males	8.7 \pm 1.0	8.7 \pm 0.9	3.7 \pm 0.6	3.7 \pm 0.4	5.9 \pm 0.6	3.5 \pm 0.6	17.2 \pm 2.8	16.8 \pm 2.5	14.7 \pm 2.4	23.7 \pm 2.6	2.0 \pm 0.5
<i>N</i> = 15	(7.2–11.0)	(7.2–10.1)	(2.7–5.0)	(3.1–4.4)	(4.2–7.2)	(2.3–4.6)	(14.4–24.6)	(13.9–23.3)	(11.2–18.8)	(19.2–27.6)	(0.9–2.7)
Females	10.4 \pm 0.6	10.5 \pm 1.4	4.7 \pm 0.7	4.3 \pm 0.6	7.2 \pm 0.5	4.6 \pm 0.7	19.7 \pm 1.0	19.6 \pm 1.6	16.7 \pm 0.5	27.0 \pm 0.9	2.9 \pm 0.2
<i>N</i> = 3	(9.8–10.9)	(9.0–11.9)	(3.9–5.3)	(3.8–4.9)	(6.7–7.7)	(4.1–5.4)	(18.7–20.7)	(17.8–20.6)	(16.3–17.2)	(26.0–27.9)	(2.6–3.0)
<i>Ophryophryne elfina</i> sp. n.											
Males	7.3 \pm 0.8	7.0 \pm 0.9	2.6 \pm 0.5	2.9 \pm 0.4	4.8 \pm 0.5	3.0 \pm 0.6	14.0 \pm 1.7	13.7 \pm 1.3	11.9 \pm 1.0	19.8 \pm 2.2	1.6 \pm 0.4
<i>N</i> = 29	(6.1–8.9)	(4.8–9.4)	(1.1–3.5)	(2.0–4.1)	(3.7–6.0)	(1.4–4.1)	(11.7–19.6)	(11.3–16.1)	(10.4–14.7)	(17.1–28.0)	(1.2–2.5)
Females	9.0 \pm 1.1	8.8 \pm 0.7	3.7 \pm 0.7	3.9 \pm 0.5	5.8 \pm 0.8	4.1 \pm 0.9	17.1 \pm 2.0	16.3 \pm 1.0	14.9 \pm 0.7	24.3 \pm 2.1	2.0 \pm 0.3
<i>N</i> = 6	(7.7–10.1)	(7.7–9.7)	(2.9–4.5)	(3.3–4.7)	(4.9–7.0)	(3.1–5.2)	(14.9–19.8)	(15.5–17.8)	(13.9–15.7)	(21.8–27.2)	(1.5–2.4)
<i>Ophryophryne synoria</i>											
Males	12.6 \pm 1.3	11.0 \pm 1.1	4.6 \pm 0.6	4.7 \pm 0.6	7.5 \pm 0.6	4.5 \pm 0.6	20.6 \pm 1.6	20.7 \pm 2.2	18.4 \pm 2.0	29.3 \pm 2.8	2.5 \pm 0.4
<i>N</i> = 14	(10.4–14.7)	(9.2–11.1)	(3.7–5.8)	(3.7–5.6)	(6.2–8.4)	(3.6–5.5)	(17.7–23.3)	(18.1–25.1)	(15.0–22.4)	(24.0–33.5)	(1.9–3.1)
Females	14.3 \pm 1.0	13.7 \pm 3.3	5.1 \pm 0.5	5.9 \pm 1.0	8.4 \pm 0.9	5.1 \pm 1.4	23.4 \pm 2.0	24.3 \pm 1.5	20.7 \pm 4.6	32.3 \pm 5.0	3.2 \pm 0.4
<i>N</i> = 3	(13.6–15.5)	(11.0–17.4)	(4.7–5.7)	(5.1–7.0)	(7.6–9.3)	(4.3–6.7)	(21.3–25.3)	(22.7–25.7)	(15.7–24.8)	(26.8–36.4)	(2.8–3.6)




Figure 3. Boxplots of SVL showing body size variation among adult *Ophryophryne* males and females from the Langbian Plateau. Horizontal lines within each box represent the median, and boxes encompass the 75th and 25th quartiles. Color of boxes/*Ophryophryne* species corresponds to icon colors on Figs 1, 2.

in Fig. 5. Advertisement calls of all three *Ophryophryne* species found on the Langbian Plateau represent a series of high whistling sounds, resembling vocalizations of passerine birds. All males vocalized in the evening from dusk until late night. The males of the undescribed *Ophryophryne* species called from stone walls above small streams or from fern leaves, the males of *Ophryophryne gerti* usually called while sitting on bush branches and fern leaves above small streams, and the males of *Ophryophryne synoria* called from stones on the rocky stream banks.

Advertisement calls of all studied species were similarly uttered in series (Fig. 5). The call repetition rate/recording/series were one of the most significant differentiating parameters between the three species (Table 4). Parameters values varied both between and within recordings for each of the three species. Though some parameters values overlapped between species, the means of most of the parameters differed significantly.




Figure 4. Two-dimensional plots of **A** the first two factors of PCA, and **B** the first and third factors of morphological characters for the *Ophryophryne* specimens examined. Data are given for males only. Star indicates the holotype specimen for each species. Circle color corresponds to those given in Fig. 1.

The frequency of maximum amplitude always coincided with the fundamental frequency and greatly varied within recordings: from 4030 to 4920 Hz for the undescribed *Ophryophryne* species, from 4450 to 5100 Hz for *O. gerti*, and from 3600 to 3890 Hz for *O. synoria*. The values of the maximum amplitude frequency in *O. synoria* were the lowest and least variable among the three species. The minimum fundamental frequency always coincided with the initial fundamental frequency whereas the maximum fundamental frequency either could coincide with the final fundamental frequency, or was close to it. Thus, the frequency modulation was expressed either in lift of fundamental frequency during the whole call, or in an unsymmetrical arch with the peak shifted to the end of the frequency band. The form of frequency modulation varied between these two forms for each species but arch-formed calls appeared most of time in *O. synoria*. The frequency range expressing depth of frequency modulation also varied within each species' calls: from 40 to 840 Hz for the undescribed *Ophryophryne* species, from 370 to 890 Hz for *O. gerti* and from 230 to 650 Hz for *O. synoria* (Table 4).

Number of harmonics varied between/within recordings but this characteristic mostly depended on recording quality (e.g., sensitivity of recording equipment, distance from vocalizing animal, signal volume and background noise). Calls from the highest quality recordings (of the undescribed *Ophryophryne* species) contained two harmonics but a major portion of other calls contained only one harmonic.

We had two sets of the undescribed *Ophryophryne* species recordings which were made at different temperatures (11.3–11.4°C in February and 17.0–17.5°C in April). Values of frequency call parameters didn't significantly differ between the two sets. However, statistically significant differences were found in several temporal parameters of the calls (data summarized in Appendix 3). For instance, the number of calls per series was 10.53 ± 0.72 (3–21, $N = 47$) in February vs. 14 ± 0.45 (2–22, $N = 93$) in April ($F_{1,138} = 18.2$, $p < 0.001$, one-way ANOVA), the inter-call duration comprised 271 ± 4.92 ms (102–942 ms, $N = 448$) in February vs. 184 ± 1.72 (96–621 ms, $N = 1209$) in April ($H_{1,1657} = 349.2$, $p < 0.001$, Kruskal-Wallis ANOVA), the call repetition rate per series was 3.18 ± 0.1 calls/s (1.33–4.91 calls/s, $N = 47$) in February vs. 4.22 ± 0.08 calls/s (2.92–5.49 calls/s, $N = 93$) in April ($F_{1,138} = 63.9$, $p < 0.001$, one-way ANOVA), call duration was also significantly longer in February 79 ± 0.32 ms (48–102 ms, $N = 496$) than in April 70 ± 0.27 ms (25–112 ms, $N = 1301$) ($H_{1,1798} = 380.2$, $p < 0.001$, Kruskal-Wallis ANOVA). April and February recordings showed no significant differences in series duration, inter-series interval duration, and call repetition rate per recording.

Taxonomic implications

Our study, based on three lines of evidence — phylogenetic analysis and distribution of mtDNA haplotypes (Figs 1, 2), multivariate statistical analysis of 23 standard morphometric traits (Fig. 4), and acoustic analysis of advertisement calls (Fig. 5), strongly indicate the presence of three independent and distinct evolutionary lineages of *Ophryophryne* on

Table 4. Measurements of advertisement call parameters for three species of *Ophryophryne* from the Langbian Plateau, and one-way ANOVA/Kruskal-Wallis results for comparison (* $p < 0.001$) between *Ophryophryne elfina* sp. n., *O. gerti* and *O. synoria*. Parameter values are given as means \pm SE (min–max). Abbreviations: *N* — number of recordings/series/calls, *F* — frequency, *s* — seconds, *ms* — milliseconds, *Hz* — hertz (continues on next page).

Parameters	<i>O. elfina</i> sp. n.	<i>O. gerti</i>	Tukey/ Mann-Whitney U post hoc tests	<i>O. synoria</i>	Tukey/ Mann-Whitney U post hoc tests	ANOVA/ Kruskal-Wallis results
Number of males	3	3	—	3	—	—
Number of recordings	5	3	—	3	—	—
Number of series	140	115	—	15	—	—
Number of calls	1797	533	—	200	—	—
Call repetition rate per recording (calls/s)	1.18 \pm 0.2 (0.77–1.95) <i>N</i> = 5	0.35 \pm 0.14 (0.11–0.59) <i>N</i> = 3	$p < 0.05$	3.07 \pm 0.13 (2.82–3.24) <i>N</i> = 3	$p < 0.001$	$F_{2,8} = 46.7^*$
Number of calls per series	12.84 \pm 0.41 (2–22) <i>N</i> = 140	4.64 \pm 0.16 (1–8) <i>N</i> = 115	$p < 0.001$	13.33 \pm 1.4 (3–24) <i>N</i> = 15	$p < 0.05$	$F_{2,267} = 151.4^*$
Series duration (s)	3.42 \pm 0.11 (0.43–9.00) <i>N</i> = 140	2.18 \pm 0.09 (0.57–8.31) <i>N</i> = 108	$p < 0.001$	2.59 \pm 0.33 (0.62–5.21) <i>N</i> = 15	$p = 0.88$	$F_{2,267} = 40.4^*$
Call repetition rate per series (calls/s)	3.87 \pm 0.07 (1.33–5.49) <i>N</i> = 140	2.33 \pm 0.03 (0.96–3.53) <i>N</i> = 108	$p < 0.001$	5.34 \pm 0.15 (4.4–6.59) <i>N</i> = 15	$p < 0.001$	$F_{2,260} = 220.7^*$
Call duration (ms)	73 \pm 0.23 (25–112) <i>N</i> = 1797	104 \pm 0.56 (75–152) <i>N</i> = 533	$p < 0.001$ (<i>U</i> = 12535.5)	62 \pm 0.46 (37–85) <i>N</i> = 200	$p < 0.001$ (<i>U</i> = 64227.5)	$H_{2,2530} = 1345.1^*$
Inter-calls interval (ms)	207 \pm 2.06 (96–942) <i>N</i> = 1657	421.54 \pm 4.17 (275–813) <i>N</i> = 418	$p < 0.001$ (<i>U</i> = 24725.5)	143 \pm 3.32 (56–528) <i>N</i> = 185	$p < 0.001$ (<i>U</i> = 64860)	$H_{2,2260} = 1008.5^*$
Inter-series interval (s)	6.51 \pm 0.41 (1.26–31.65) <i>N</i> = 135	7.98 \pm 0.55 (0.65–39.76) <i>N</i> = 112	$p < 0.001$ (<i>U</i> = 5593.5)	1.64 \pm 0.19 (0.88–2.83) <i>N</i> = 12	$p < 0.001$ (<i>U</i> = 51)	$H_{2,2359} = 42.7^*$

Parameters	<i>O. elfina</i> sp. n.	<i>O. gerti</i>	Tukey/ Mann-Whitney U post hoc tests	<i>O. synoria</i>	Tukey/ Mann-Whitney U post hoc tests	ANOVA/ Kruskal-Wallis results
F initial (Hz)	4348.02 ± 2.96 (3980–4680) <i>N</i> = 1797	4414.17 ± 5.12 (4070–4640) <i>N</i> = 533	p < 0.001 (U = 332858)	3449.55 ± 6.41 (3230–3700) <i>N</i> = 200	p < 0.001 (U = 0)	H _{2,2530} = 655.7*
F final (Hz)	4715.3 ± 3.29 (4260–5010) <i>N</i> = 1797	4888.76 ± 3.82 (4640–5150) <i>N</i> = 533	p < 0.001 (U = 142738)	3708.9 ± 9.28 (3420–3980) <i>N</i> = 200	p < 0.001 (U = 0)	H _{2,2530} = 1075.9*
F maximum (Hz)	4807.74 ± 3.46 (4260–5060) <i>N</i> = 1797	4998.74 ± 4.27 (4780–5250) <i>N</i> = 533	p < 0.001 (U = 138045)	3907.05 ± 4.22 (3750–4070) <i>N</i> = 200	p < 0.001 (U = 0)	H _{2,2530} = 1094.2*
F minimum (Hz)	4348.02 ± 2.96 (3980–4680) <i>N</i> = 1797	4414.17 ± 5.12 (4070–4640) <i>N</i> = 533	p < 0.001 (U = 309752.5)	3449.55 ± 6.41 (3230–3700) <i>N</i> = 200	p < 0.001 (U = 0)	H _{2,2530} = 689.5*
F range (Hz)	459.71 ± 3.27 (40–840) <i>N</i> = 1797	584.58 ± 3.87 (370–890) <i>N</i> = 533	p < 0.001 (U = 229934)	457.5 ± 5.86 (230–650) <i>N</i> = 200	p = 0.61 (U = 175721)	H _{2,2530} = 367.1*
F peak (Hz)	4645.94 ± 4.39 (4030–4920) <i>N</i> = 1797	4845.99 ± 4.22 (4450–5100) <i>N</i> = 533	p < 0.001 (U = 157981.5)	3798.9 ± 4.87 (3600–3890) <i>N</i> = 200	p < 0.001 (U = 0)	H _{2,2530} = 1030.2*




Figure 5. Oscillograms (top) and sonograms (bottom) of male advertisement calls of **A** *Ophryophryne elfina* sp. n. (Lam Dong Prov., Bidoup-Nui Ba N.P., 1935 m a.s.l., 17°C) **B** *O. gerti* (Dak Lak Prov., Chu Yang Sin N.P., 1020 m a.s.l., 22°C) **C** *O. synoria* (Dak Lak Prov., Chu Yang Sin N.P., 750 m a.s.l., 21°C). The sampling rate lowered to 22.05 kHz.

the Langbian Plateau and in surrounding areas of southern Truong Son Mountains in southern Vietnam and eastern Cambodia. Our examination of type material allowed us to identify the two larger lineages as *O. synoria* and *O. gerti*, whereas the smaller species of *Ophryophryne* represents an undescribed taxon (Fig. 6). In two of the 12 surveyed localities (see Fig. 1) all three species were recorded in sympatry, with *O. synoria* and *O. gerti* recorded syntopically in the same streams in environs of Giang Ly Ranger Station, Bidoup–Nui Ba N.P., Lam Dong Prov. (Loc. 6, Fig. 1) and in Chu Yang Sin N.P. in Dak Lak Prov. (Loc. 11, Fig. 1); the small-sized undescribed *Ophryophryne* species was usually found in higher elevations, but was nevertheless recorded synbiotically with *O. gerti*




Figure 6. *Ophryophryne* species of the Langbian Plateau (Vietnam): **A** *O. gerti*, female, Chu Yang Sin N.P., Dak Lak Prov., 1000 m a.s.l. **B** *O. gerti*, male, Chu Yang Sin N.P., Dak Lak Prov., 1000 m a.s.l. **C** *O. gerti*, male, Bidoup–Nui Ba N.P., Lam Dong Prov., 1550 m a.s.l. **D** *O. synoria*, female, Bu Gia Map N.P., Binh Phuoc Prov., 400 m a.s.l. **E** *O. synoria*, male, Chu Yang Sin Mt., Chu Yang Sin N.P., Dak Lak Prov., 1000 m a.s.l. **F** *O. synoria*, male, Bidoup–Nui Ba N.P., Lam Dong Prov., 1550 m a.s.l. **G** *Ophryophryne elfina* sp. n., male, Chu Yang Sin Mt., Chu Yang Sin N.P., Dak Lak Prov., 2000 m a.s.l. **H** *Ophryophryne elfina* sp. n., male, Hon Giao Mt., Bidoup–Nui Ba N.P., Lam Dong and Khanh Hoa provincial border, 2000 m a.s.l. Photos by N.A. Poyarkov and N.L. Orlov.

in Chu Yang Sin N.P. in Dak Lak Prov. (Loc. 11, Fig. 1). Below we provide taxonomic remarks on *Ophryophryne* species of the Langbian Plateau along with the description of the new species.

Taxonomic remarks on *Ophryophryne gerti* Ohler, 2003

Chresonymy:

[?] *Ophryophryne microstoma* — Orlov et al. 2002:84 (partim—Dac Lac [Dak Lak] and Lam Dong provinces, Vietnam; no referred specimens indicated, inclusion *fide* N.L. Orlov).

Ophryophryne gerti Ohler 2003:25, fig. 1 (partim—BMNH 1921.4.1.324, BMNH 1921.4.1.323).

[?] *Ophryophryne gerti* — Ohler 2003:25; Stuart et al., 2006:135 (FMNH 252899, FMNH 252901).

[?] *Ophryophryne microstoma* — Nguyen et al. 2005:15 (partim—“Dak Lak and Lam Dong” provinces, Vietnam; following indication by Orlov et al. 2002).

“*Ophryophryne* sp. 2” — Orlov et al. 2008:82 (Chu Yang Sin N.P., Dak Lak Prov., Vietnam; ZISP 12836–12879).

Ophryophryne gerti — Nguyen et al. 2009:84–85 (partim—“Lam Dong (Cam Li, Dran)”, Vietnam).

[?] *Ophryophryne microstoma* — Nguyen et al. 2009:86 (partim—“Dak Lak, Lam Dong”, Vietnam; based on the record by Orlov et al. 2002).

[?] *Ophryophryne gerti* — Stuart et al. 2010:40 (eight uncataloged “topotype” females from Bidoup–Nui Ba N.P., Langbian Plateau, Vietnam).

Ophryophryne gerti — Poyarkov [Paiarkov] and Vassilieva 2011:202 (Bidoup–Nui Ba N.P., Lam Dong Prov., Vietnam; ZMMU A-4715, ZMMU A-4718).

Removed from chresonymy:

Ophryophryne gerti Ohler 2003:25 (partim—BMNH “1972.15.2.4” [sic. BMNH 1972.1524])

Ophryophryne gerti — Stuart 2005:475 (FMNH 258564)

Ophryophryne gerti — Bain et al. 2007:108 (AMNH A-169287, AMNH A-163668)

[?] *Ophryophryne gerti* — Orlov et al. 2015:2010, fig. 9 (Gia Lai Prov., Vietnam; based on identification of the specimen illustrated in fig. 9 as *Ophryophryne* cf. *poilani* *sensu auctorum*).

Holotype. BMNH 1921.4.1.324, adult male from “Cam Ly (river), south-east of Da Lat (11°56'N; 108°25'E), Lang Bian Plateau, sLam [sic. = Lam] Dong Province, Vietnam”, collected by M.A. Smith, presented to BMNH in 1921 (Ohler 2003; NHMUK specimen catalogue). Re-examined by SM.

Paratypes. BMNH 1921.4.1.323, immature female from “Dran (11°50'N; 108°34'E), Lang Bian Plateau, Lam Dong Province, Vietnam”, collected by M.A.

Smith, presented to BMNH in 1921; BMNH 1972.1524 (see ‘Remarks’) adult male, “Huey Sapan, Pak Maat (precise location not found), Mekong, Laos”, collector M.A. Smith, accessioned in the BMNH from Smith’s private collection in 1972, collection date unknown (Ohler 2003; NHMUK specimen catalogue).

Remarks. Ohler (2003) provides the paratype number BMNH 1972.15.2.4 for the Laos paratype specimen, however, the number on the specimen tag and NHMUK specimen catalogue reads BMNH 1972.1524 (Mahony 2011b). Our present data suggests that *O. gerti* is not found beyond the limits of the Langbian Plateau, thus we are confident that the paratype BMNH 1972.1524 does not represent this biological taxon. Ohler (2003: and by implication Stuart et al. 2006) preliminarily identified two specimens from Ankhe Dist. in northern Gia Lai Prov. as *O. gerti*. This locality is disconnected from the Langbian highlands by a wide lowland area, indicating that these specimens are biogeographically isolated from *O. gerti s. stricto*. Further work is necessary to ascertain the taxonomic status of these specimens. Orlov et al. (2002), without providing data on examined specimens, considered that the distribution of *O. microstoma* extended south in Vietnam as far as the Lam Dong and Dak Lak provinces. Our results suggest that these southern Vietnamese populations most likely represented the superficially similar *O. gerti*, *O. synoria*, or possibly the new taxon described below. Stuart (2005) reports *O. gerti* from Champasak Prov. in southern Laos, based on a single specimen (FMNH 258564: not examined here). This locality is biogeographically not connected to the known range of this species, as redefined here, thus the taxonomic status of this specimen requires further confirmation. Bain et al. (2007) identifies two specimens AMNH A-169287 (Thua Thien-Hue Prov., Vietnam) and AMNH A-163668 (Quang Nam Prov., Vietnam) as *O. gerti*. We re-examined these specimens and regard them to be morphologically more similar to *O. poilani*. Furthermore, both specimens were included in our molecular analysis and found to be distantly related to *O. gerti s. stricto* (Fig. 2). Stuart et al. (2010) provides the SVL range (SVL 37.5–42.5 mm, mean \pm SD 40.4 \pm 1.6, $N = 8$) for ‘topotype’ female specimens of *O. “gerti”*, however these specimens are smaller than females of *O. gerti* provided herein (SVL 43.1–47.4 mm, mean \pm SD 45.07 \pm 2.16, $N = 3$), and larger than the new taxon described below (SVL 35.1–36.5 mm, mean \pm SD 35.6 \pm 0.5, $N = 6$) (see Table 3 for details). The taxonomic status of these specimens remains unknown.

Vernacular name. English: “*Gerti’s Mountain Toad*”; Vietnamese: “*Cóc Núi Goti*” (Nguyen et al. 2009), “*Cóc Núi Goti*” (Nguyen et al. 2014).

Redescription of the holotype. Mature male (SVL 35.7 mm), habitus slender (Fig. 7A, B). Specimen in good state of preservation; two incisions are present on trunk, one longitudinally orientated on mid-abdomen, another longitudinally orientated on upper flank on right side; liver and testes observable through incisions, testes enlarged; jaw is dislocated on right allowing visual access to buccal cavity.

Head moderately small (HL/SVL 25.5%; HW/SVL 29.7%), wider than long (HW/HL 116.5%), relatively deep; top of head flat; snout comparatively short (SL/HL 33.0%), truncated in dorsal view (Fig. 7A), projecting significantly beyond margin of lower jaw in profile (Fig. 7C); loreal region vertical; canthus rostralis distinct, moderately sharp; dorsal region of snout flat; eyes large (ED/HL 38.5%), slightly protuberant in dorsal view and in profile, horizontal eye diameter less than twice as long

as maximum tympanum diameter (TYD/ED 62.9%) and longer than snout (ED/SL 116.7%); eye to tympanum distance approximately equal to maximum tympanum diameter (TYE/TYD 95.5%); tympanum distinct, circular, moderately large (TYD/HL 24.2%); pupil in preservation oval (Fig. 7C), vertically orientated; nostril opening oval, vertical, laterally orientated, medially located between eye and snout (EN/NS 100.0%); internarial distance subequal to upper eyelid width (IN/UEW 103.2%), and significantly less than narrowest point between upper eyelids (IN/IUE 145.5%); pineal ocellus not visible externally; vomerine ridges absent; maxillary and vomerine teeth absent; vocal sac gular, its' openings not discernable; tongue moderately large, posterior end free, with weak notch posteriorly, lacking medial lingual process.

Forelimbs slender, forearm moderately long (FAL/SVL 22.4%) slightly enlarged relative to upper forelimb, and shorter than hand (FAL/HAL 95.2%); fingers long and narrow, dorsoventrally flattened; lateral fringes and webbing absent (Fig. 7D); finger length formula $F1 = F2 < F4 < F3$; subarticular tubercles absent, replaced by low callous dermal ridges; supernumerary tubercles absent; outer palmar (metacarpal) tubercle weak, longitudinally oval, elevated but with indistinct borders (Fig. 7D); thenar tubercles weak; finger tips rounded, weakly expanded relative to digit widths (wider than distal-most finger articulation), with circular pads (Fig. 7D); terminal grooves absent.

Hindlimbs slender, relatively long, shanks overlap when thighs are held at right angle to body; shank length less than half of snout to vent length (SHL/SVL 44.5%); thighs shorter than shanks (SHL/TL 106.0%), and feet (FOL/TL 106.4%); toes long and slightly dorsoventrally flattened (Fig. 7E), relative toe lengths $T1 < T2 < T5 < T3 < T4$; lateral fringes on toes, outer metatarsal tubercle, subarticular and supernumerary tubercles absent; inner metatarsal tubercle well developed with distinct borders, oval-shaped (IMT/FOL 11.3%) (Fig. 7E); weak ridge of callous tissue present on ventral surface of all toes, not continuing onto metatarsus; webbing between digits rudimentary; tarsal fold absent; toe tips not expanded relative to digit widths, with circular pads; terminal grooves absent.

Skin texture and skin glands in preservation. Skin of dorsal and lateral surfaces of head, body and limbs smooth with numerous small tubercles finely and relatively evenly scattered on dorsal surfaces of trunk, head and limbs (Fig. 7A); small tubercles present on temporal region, tympanum smooth with borders weakly raised; tubercles arranged in distinct transverse ridges on dorsal surfaces of forearms, shanks and thighs; numerous large tubercles on flanks irregularly scattered from axilla to groin, intermixed with smaller tubercles; central portion of outer edge of upper eyelids slightly thickened, with a single short tubercular spine (Fig. 7C), transverse fold on posterior edges of upper eyelids absent; well-developed glandular supratympanic folds, narrow anteriorly, considerably widening posteriorly, from posterior corner of orbits, extending along upper margin of tympanum, terminating above forelimb insertions (Fig. 7C); dorsolateral glandular ridge well-developed, extending from posterior to supratympanic ridges to ca. 75% of trunk length, on each side; a moderately well-developed “>-<” shaped glandular parietoscapular-sacral ridge present on dorsum (see Fig. 7A); two small tubercles present above vent; gular




Figure 7. Holotype *Ophryophryne gerti* (BMNH 1921.4.1.324, male) in preservative: **A** dorsal view **B** ventral view **C** head, lateral view **D** volar view of the left hand **E** plantar view of the left foot. Photos by S. Mahony.

region, chest, and ventral surfaces of limbs smooth to weakly shagreened, abdomen weakly granular; two nuptial pads per limb, one large on dorsal surface of F1 from base of metacarpal to near distal joint, another small pad on inner dorsal surface of F2 on metacarpal; pectoral glands round, weakly raised, positioned level with axilla; femoral glands slightly raised, average size, on posterior surface of thighs, situated slightly closer to knee than to cloaca; numerous small white asperities present on posterior half of dorsum, sparse anteriorly, increasing in density posteriorly to above cloaca, absent from all remaining surfaces.

Color of holotype in preservative (Figure 7). Dorsal surfaces of head, body, forearms and hindlimbs mid to light brown, slightly lighter on flanks and dorsal surface of upper arms; a distinct darker brown “V”-shaped marking on dorsal surface of head; no distinct “X”-shaped or hourglass marking on mid dorsum; most flank tubercles are bordered below by a small dark brown spot anteriorly, increasing in size posteriorly towards groin; a broad brown stripe extends around lateral surfaces of snout, from anterior borders of orbits between canthus rostralis and the level of lower border of nostrils; two broad darker brown vertical stripes below orbits, one at level of anterior orbital border, and a second extends from central lower border of orbits to edge of jaw, a faint darker brown stripe extends from posterior border of orbits to cover tympanum; color of supratympanic folds same as surrounding surfaces, but lower border dark brown; edge of lower eyelid dark brown; dorsal surfaces of forearms each with two dark brown transverse blotches, and thighs and shanks with faint darker brown transverse stripes; ventral surfaces of throat, chest and anterior half of abdomen, and ventral surface of hands primarily plain light brown, fading to a mottled brownish beige with small dark brown blotches on posterior half of abdomen, and ventral surfaces of forelimbs, thighs and shanks, and dorsal surfaces of tarsi and feet; ventral surface of tarsi and feet dark brown fading distally on toes to a mid-brown; area surrounding cloaca dark brown, fading distally on lateral surfaces of thighs.

Measurements of the holotype (all in mm, taken by SM). SVL 35.7; HW 10.6; HL 9.1; IFE 5.0; IBE 8.2; ED 3.5; TYD 2.2; TYE 2.1; SL 3.0; EN 1.5; NS 1.5; IUE 2.2; IN 3.2; UEW 3.1; FAL 8.0; HAL 8.4; FIL 3.4; FIIL 3.4; FIIL 5.6; FIVL 3.6; SHL 15.9; TL 15.0; FOL 14.1; TFOL 21.7; IMT 1.6; TIL 1.7; TIIL 3.8; TIIL 5.5; TIVL 6.7; TVL 3.5.

Distribution. *Ophryophryne gerti* is herein confirmed from three localities on the Langbian Plateau in southern Vietnam, between 700–2000 m a.s.l. (Fig. 1): Cam Ly River and Nui Ba Mt. in environs of Dalat city, Lam Dong Prov., Vietnam (1000–1800 m a.s.l.) (Ohler 2003, this study); Environs of Bidoup Mt. (2000 m a.s.l.), and Giang Ly Ranger Station (1500 m a.s.l.), Bidoup–Nui Ba N.P., Lam Dong Prov., Vietnam (this study); Chu Yang Sin Mt. environs, Krong Kmar Commune, Krong Bong Dist., Dak Lak Prov., Vietnam (700–2000 m a.s.l.) (Orlov et al. 2008; this study). Additional localities reported in literature require confirmation pending further study of voucher material (see Remarks section above).

New records and range extension for *Ophryophryne synoria* Stuart, Sok & Neang, 2006

Chresonymy:

Ophryophryne synoria Stuart et al. 2006:135, fig. 5.

“*Ophryophryne* sp. 3” — Orlov et al. 2008:82 (Chu Yang Sin N.P., Dak Lak Prov., Vietnam; ZISP 12811–12833).

[?] *Ophryophryne* sp. cf. *poilani* — Poyarkov [Paiarkov] and Vassilieva 2011:202 (Bidoup Mt., Bidoup–Nui Ba N.P., Lam Dong Prov., Vietnam; ZMMU A-4713).

Ophryophryne synoria — Vassilieva et al. 2016:54–56, figs. 44–47 (Binh Phuoc Prov. and Dong Nai Prov., Vietnam; ZMMU A-4516, ZMMU A-5003).

Holotype. FMNH 262779, adult male from “O Chung Chry Stream, near 12°17'30"N, 107°03'06"E, 500 m elev., Seima Biodiversity Conservation Area, O’Rang District, Mondolkiri Province, Cambodia”, collected by Bryan Lynn Stuart, Sok Ko and Neang Thy (Stuart et al. 2006; FMNH specimen catalogue). Re-examined by SM.

Paratype. FMNH 262778, adult male, collected with the holotype (Stuart et al. 2006).

Measurements of the holotype (all in mm, taken by SM). SVL 48.8; HW 14.2; HL 13.1; IFE 6.1; IBE 10.8; ED 4.7; TYD 3.0; TYE 3.1; SL 4.3; EN 2.5; NS 1.9; IUE 3.5; IN 3.4; UEW 4.1; FAL 12.0; HAL 13.0; FIL 5.1; FIIL 5.1; FIIIL 8.4; FIVL 5.5; SHL 21.8; TL 21.2; FOL 20.0; TFOL 29.4; IMT 2.2.

Distribution and remarks. Stuart et al. (2006) described a large-sized *Ophryophryne* from O’Rang (also spelled as “O’Reang”) District in eastern Cambodia, close to the Vietnamese border, as *O. synoria* (Loc. 1, Fig. 1). Subsequently, during field surveys in 2009–2011, the species was reported in southern Vietnam from Bu Gia Map N.P., Binh Phuoc Prov. (Loc. 2, Fig. 1) and Cat Tien N.P. in Dong Nai Prov. (Loc. 3, Fig. 1) based on morphological evidence (Vassilieva et al. 2016). Herein, we confirm the identity of these specimens based on morphological and molecular genetic evidence, and further expand its distribution in southern Vietnam to include medium and low elevation localities in the central and western parts of the Langbian Plateau (Dak Lak, Lam Dong, Dong Nai and Binh Phuoc provinces between 200 and 1500 m a.s.l.; its presence in Dak Nong Prov. is anticipated). We also identify two mtDNA lineages within *O. synoria* with a moderate level of sequence divergence ($p = 2.6\%$: Table 2, Fig. 2): Subclade B inhabits mountain areas in Lam Dong and Dak Lak provinces and was also recorded for the lowland habitat in Dong Nai Prov. (Locs. 3–4, 6 and 11, Fig. 1) whereas Subclade C is only found in Mondolkiri Prov. of Cambodia and adjacent Binh Phuoc Prov. of Vietnam (Fig. 1, Locs 1–2) and corresponds to *O. synoria s. stricto*.

Variation. The studied specimens of *O. synoria* showed substantial variation in morphological characters, including SVL (Fig. 3) and other morphometric characters (Fig. 4), coloration, and degree of development of palpebral projection (Fig. 6D, E and F). Overall morphology, coloration, and skin glands of the newly discovered populations of *O. synoria* are in general agreement with the description of the holotype by Stuart et al. (2006). Young specimens from Bidoup–Nui Ba N.P. in life often

have reddish or orange coloration of thighs and groin, which was not observed in the type specimens from Cambodia (Stuart et al. 2006), nor in the Bu Gia Map population. The degree of development of short dorsolateral glandular folds varied among specimens, but they were always distinct (Fig. 6D, E and F). The holotype and the Bu Gia Map population (Subclade C) have the finger length formula $F1 = F2 < F4 < F3$, while in populations from Bidoup–Nui Ba N.P. and Chu Yang Sin N.P. (Subclade B), the finger length formula is $F1 = F4 < F2 < F3$. Subclade B populations also tend to have a slightly larger tympanum (TYD/ED 68.4%–80.1%; TYE/TD 73.3%–80.6%) than the nominative *O. synoria* (TYD/ED 62.0%–71.9%; TYE/TD 83.9%–103.3%). Though the taxonomic status of the two revealed lineages is not completely clear, here-in we tentatively regard them as deep intraspecific mtDNA lineages based on observed genetic differentiation and overall morphological similarity.

Vernacular name. English: “*O’Reang Mountain Toad*” (this paper); “*O’Reang horned frog*” (Vassilieva et al. 2016); Vietnamese: “*Cóc Núi O-Reng*” (Vassilieva et al. 2016).

Description of a new species of *Ophryophryne*

Based upon several lines of evidence, including the analyses of diagnostic morphological characters, acoustic analyses of advertisement calls and phylogenetic analyses of mtDNA sequences for the 12S rRNA–16S rRNA genes, the new species of *Ophryophryne* from mid to high elevations of the western Langbian Plateau represents a highly divergent mtDNA lineage, clearly distinct from all other *Ophryophryne* species. These results support our hypothesis that this recently discovered lineage of *Ophryophryne* represents an undescribed species, described below:

Ophryophryne elfina sp. n.

<http://zoobank.org/481B0CFA-5428-40E1-A7A3-C132DC840EC0>

Figs 6G, H, 8–12, 13C, 14A (right), 14B, C

Chresonymy:

“*Ophryophryne* sp. 1” — Orlov et al. 2008:82 (Chu Yang Sin N.P., Dak Lak Prov., Vietnam; ZISP 12880–12884).

[?] *Ophryophryne gerti* — Stuart et al. 2010:40 (eight uncataloged topotype females from Bidoup–Nui Ba N.P., Langbian Plateau, Vietnam).

“*Ophryophryne* sp.” — Poyarkov [Paiarkov] and Vassilieva 2011:174, 202; fig. 5.6 (Bidoup Mt., Bidoup–Nui Ba N.P., Lam Dong Prov., Vietnam; ZMMU A-4716, ZMMU A-4788, ZMMU A-5674, ZMMU A-5675).

Ophryophryne gerti — Nguyen et al. 2014:148–149; fig. 2 (partim—Hon Ba Mt., Hon Ba Nature Reserve [hereafter N.R.], Khanh Hoa Prov., Vietnam; VNMN 983, ZFMK 94220).

Holotype. ZMMU A-5669 (field number NAP-02658), adult male from the northern slope of Chu Pan Fan Mountain, Chu Yang Sin National Park, Bong Krang Commune, Lak District, Dak Lak Province, Vietnam (coordinates 12°22'31.90"N; 108°21'14.10"E, elevation 1725 m a.s.l.), collected along a mountain stream in mixed evergreen montane tropical forest by N.A. Poyarkov on 07 April 2012 (Figs 8 and 9).

Paratypes. ZMMU A-5691 (field numbers ABV-00580; ABV-00581), two juveniles from the north-western slope of Chu Yang Sin Mountain, Chu Yang Sin N.P., Hoa Le Commune, Krong Bong Dist., Dak Lak Prov., Vietnam (12°24'47.70"N; 108°24'25.40"E, 1975 m a.s.l.), collected in leaf litter in mixed evergreen montane tropical forest by N.A. Poyarkov on 25 May 2014; ZMMU A-5692 (field number NAP-00582), adult male from the northern slope of Bidoup Mountain, Bidoup–Nui Ba N.P., Da Chais Commune, Lac Duong Dist., Lam Dong Prov., Vietnam (12°06'5.60"N; 108°39'34.20"E, 2035 m a.s.l.), collected along mountain stream in evergreen montane elfin forest by N.A. Poyarkov on 09 May 2009; ZMMU A-5675 (field numbers NAP-01456; NAP-01459), two adult males from the northern slope of Bidoup Mountain, Bidoup–Nui Ba N.P., Da Chais Commune, Lac Duong Dist., Lam Dong Prov., Vietnam (12°06'5.60"N; 108°39'34.20"E, 2035 m a.s.l.), collected on stones and vegetation along a mountain stream in mixed evergreen montane elfin forest by N.A. Poyarkov and A.B. Vassilieva on 25 June 2010; ZMMU A-4788 (field numbers NAP-01455; NAP-01449; NAP-01450; NAP-01460), four adult males from the southern slope of Hon Giao Mountain Ridge, Bidoup–Nui Ba N.P., Da Chais Commune, Lac Duong Dist., border of Lam Dong Prov. and Khanh Hoa Prov., Vietnam (12°11'33.10"N; 108°42'41.80"E, 1890 m a.s.l.), collected along a mountain stream sitting on stones and tree branches near the water edge, in mixed evergreen montane elfin forest by N.A. Poyarkov and A.B. Vassilieva on 30 June 2010; ZMMU A-5674 (field numbers NAP-01451; NAP-01452), two adult males from the southern slope of Hon Giao Mountain Ridge, Bidoup–Nui Ba N.P., Da Chais Commune, Lac Duong Dist., border of Lam Dong Prov. and Khanh Hoa Prov., Vietnam (12°11'33.10"N; 108°42'41.80"E, 1890 m a.s.l.), collected along a mountain stream on stones and vegetation in evergreen montane elfin forest by N.A. Poyarkov and A.B. Vassilieva on 29 June 2010; ZMMU A-5170 three adult males (field numbers ABV-00454; ABV-00472; ABV-00471), and one adult female (ABV-00455) from the northern slope of Bidoup Mountain, Bidoup–Nui Ba N.P., Da Chais Commune, Lac Duong Dist., Lam Dong Prov., Vietnam (12°06'5.60"N; 108°39'34.20"E, 2035 m a.s.l.), collected along a mountain stream in mixed evergreen montane tropical forest by A.B. Vassilieva on 16 April 2014.

Referred specimens. ITBCZ 2786, ITBCZ 2788, ITBCZ 2792, ITBCZ 2828, three adult females and one adult male collected along a mountain stream in evergreen mountain forest on the summit of Hon Ba Mountain, Hon Ba N.R., Dien Khanh Dist., Khanh Hoa Prov., Vietnam (12°07'10.60"N; 108°56'51.60"E, 1510 m a.s.l.), by Sang Ngoc Nguyen, Luan Thanh Nguyen and Vu Dang Hoang Nguyen on 22–24 December 2015; ITBCZ 2908–2909, ITBCZ 2918–2919, ITBCZ 3502, five adult males collected along a mountain stream in evergreen montane forest on the summit of Hon Ba Mountain, Hon Ba N.R., Dien Khanh Dist., Khanh Hoa Prov., Vietnam (12°07'28.80"N; 108°58'14.20"E, 950 m a.s.l.), by Sang Ngoc Nguyen, Luan Thanh Nguyen, and Vu

Dang Hoang Nguyen on 22–28 March 2016; ZMMU A-5679 (field number NAP-01169), 7 larvae collected in a cascade mountain stream on the northern slope of Bidoup Mountain, Bidoup–Nui Ba N.P., Da Chais Commune, Lac Duong Dist., Lam Dong Prov., Vietnam (12°06'5.60"N; 108°39'34.20"E, 2035 m a.s.l.), by N.A. Poyarkov on 03 May 2009; ZMMU A-5684 (field number NAP-02673), 4 larvae collected in a cascade mountain stream on the northern slope of Chu Pan Fan Mt., Chu Yang Sin N.P., Bong Krang Commune, Lak Dist., Dak Lak Prov., Vietnam (coordinates 12°22'31.90"N; 108°21'14.10"E, elevation 1725 m a.s.l.), by N.A. Poyarkov on 07 April 2012.

Etymology. The specific epithet is an adjective (in agreement with the genus name in feminine gender), derived from “*elf*”, the English spelling of “*alfus*” in Latin, referring to usually forest-dwelling supernatural mythological creatures in Germanic mythology and folklore; the name is given in reference both to the funny appearance and small size of the new species, as well as to the their endangered habitat, restricted to wet evergreen montane forests at high elevations of the Langbian Plateau; such forests are often called “*elfin forests*”.

Recommended vernacular name. The recommended common name in English is “Elfin Mountain Toad”. The recommended common name in Vietnamese is “Cóc Núi Tiểu Yêu Tinh”.

Diagnosis. The species is allocated to *Ophryophryne* based on its obvious similarities with its sister taxa, its molecular phylogenetic affinities, and the absence of maxillary teeth considered diagnostic for the genus (previous authors, e.g. Ohler 2003, Delorme et al. 2006 and Fei et al. 2009 also indicated a horizontal pupil and the absence of vomerine teeth as diagnostic for *Ophryophryne*, this is reconsidered by Mahony et al. 2017). *Ophryophryne elfina* sp. n. is distinguished from its congeners by a combination of the following morphological attributes: (1) small adult body size, male SVL 26.9–33.9 mm ($N = 29$), female SVL 35.1–36.5 mm ($N = 6$); (2) snout sharply protruding in profile; (3) tympanum diameter approximately half of eye diameter; tympanum to eye distance approximately 70–90% of tympanum diameter; (4) finger length formula: $F1 < F4 \leq F2 < F3$, or $F1 \leq F2 < F4 < F3$; toe webbing rudimentary, toe length formula: $T1 < T5 < T2 < T3 < T4$; (5) short dorsolateral glandular ridge present above shoulder; (6) palpebral projection present as a small single tubercle to moderately developed single projection; (7) dermal cloacal protuberance and dermal flaps above cloacal opening absent; (8) skin of dorsal and lateral surfaces of head, body and limbs shagreened with numerous small tubercles, large warts on the flanks; (9) skin on dorsal and lateral surfaces of body, head and limbs with numerous bright orange-red (in life) asperities; (10) males with a red-orange (in life) nuptial pad on F1; (11) dorsal coloration light yellow-brown with dark hourglass-shaped marking on dorsum usually edged with white or beige (in life); (12) posterior suborbital light bar well-defined, usually clearly separated from dark-brown temporal triangular spot, uniformly covering temporal area and tympanum.

The new species is also markedly distinct from all congeners for which comparable sequences are available (16S rRNA mitochondrial gene; uncorrected genetic distance $> 8.2\%$). The advertisement call of the new species consists of whistling notes uttered in series: average 12.84 ± 0.41 calls per series, with an average dominant frequency of

4645.94 \pm 4.39 Hz, repetition rate per recording/series 1.18 \pm 0.2 calls/s and 3.87 \pm 0.07 calls/s, respectively, with average call duration 73 \pm 0.23 ms and inter-call interval 207 \pm 2.06 ms, also distinguishes the new species from *Ophryophryne* species for which calls are known, including the two species found in sympatry.

Description of holotype. Mature male (SVL 27.2 mm); habitus slender (Figs 8A, B, and 9). Specimen in good state of preservation; median abdomen dissected, dissection length ca. 9.0 mm; liver and testes observable through incision, testes white, enlarged (testes length 3.9 mm; Fig. 8B); ventral right femur dissected for molecular sampling, dissection length 9.8 mm.

Head moderately large (HL/SVL 29.2%; HW/SVL 29.5%), almost as wide as long (HW/HL 101.0%), triangular in dorsal view; top of head flat; snout comparatively short (ES/HL 30.9%), narrow (IFE/HW 39.4%), somewhat truncated in dorsal view (Fig. 8A), sharply protruding in profile, without rostral appendage (Figs 8C, 9); snout projecting significantly beyond margin of lower jaw (Fig. 8C); loreal region slightly concave; canthus rostralis distinct, sharp; dorsal region of snout flat; eyes large (ED/HL 44.8%), slightly protuberant in dorsal view and in profile, eye less than twice as long as maximum tympanum diameter (TYD/ED 53.7%) and half times longer than snout (ED/SL 145.3%); eye-tympanum distance less than maximum tympanum diameter (TYE/TD 83.3%); tympanum distinct, almost circular shaped with vertical diameter slightly exceeding horizontal diameter, tympanum large (TYD/HL 24.1%); pupil in preservation diamond-shaped (Fig. 8C), horizontally orientated; nostril oval-shaped, vertical, laterally orientated, located as far from eye as from snout (EN/NS 100.8%); internarial distance greater than eyelid width (IN/UEW 119.5%), and sub-equal to narrowest point between upper eyelids (IN/IUE 103.7%); pineal ocellus not visible externally (Fig. 8A); vomerine ridges not absent; maxillary and vomerine teeth absent; vocal sac gular, its openings not discernable; tongue moderately large, with free posterior end, not notched posteriorly, lacking medial lingual process.

Forelimbs slender, forearm moderately long (FAL/SVL 25.7%), slightly enlarged relative to upper forelimb, and shorter than hand (FAL/HAL 96.1%); fingers long and narrow, dorsoventrally flattened, weak lateral fringes present on third and fourth fingers (Fig. 8D), finger length formula $F1 < F4 < F2 < F3$; fingers completely free of webbing; subarticular tubercles absent, replaced by low callous dermal ridges; supernumerary tubercles absent; outer palmar (metacarpal) tubercle small, round, elevated but with indistinct borders (Fig. 8D); thenar tubercle weak; finger tips in life rounded, weakly expanded relative to digit widths (wider than the distal-most finger articulation), with circular pads (Fig. 8D); terminal grooves absent.

Hindlimbs slender, relatively long, shanks overlap when thighs are held at right angle to body; shank length less than half of snout–vent length (SHL/SVL 48.7%); thighs shorter than shanks (SHL/TL 109.2%), and feet (FOL/TL 110.9%); toes long and slightly dorsoventrally flattened (Fig. 8E), relative toe lengths $T1 < T5 < T2 < T3 < T4$; toe tips slightly expanded relative to digit widths (wider than the distal-most toe articulation), with circular pads; terminal grooves absent; lateral fringes on toes, outer metatarsal tubercle, subarticular and supernumerary tubercles absent; inner metatarsal




Figure 8. Holotype *Ophryophryne elfina* sp. n. (ZMMU A-5669, male, field number NAP-02658) in preservative: **A** dorsal view **B** ventral view **C** head, lateral view **D** volar view of the left hand **E** plantar view of the left foot. Photos by N.A. Poyarkov.



Figure 9. Holotype of *Ophryophryne elfina* sp. n. in life (ZMMU A-5669, male, field number NAP-02658), dorsolateral view. Photos by N.A. Poyarkov.

tubercle well developed with distinct borders, ca. two times longer than wide, oval-shaped (IMT/FOL 9.3%); well-developed dermal ridge of callous tissue present on ventral surface of all toes and continuing to metatarsus; rudimentary webbing present between all five toes, basal web distinct between toes T2–T3 and T3–T4, but is completely reduced between toes T1–T2 and T4–T5; tarsal fold absent.

Skin texture and skin glands. Skin of dorsal and lateral surfaces of head, body and limbs shagreened, with numerous small skin asperities present on anterior two thirds of dorsum, sparse posteriorly, increasing in density along dermal ridges, densely covering dorsal and lateral surfaces of head, upper eyelids, and dorsal surfaces of thighs, shanks, upper forelimbs, forearms, hands, feet and digits, and absent from all remaining surfaces. Small tubercles finely and relatively evenly scattered on dorsal surfaces of trunk, head and limbs, including maxilla, mandible, eyelids and dorsal surfaces of head, forelimbs and hindlimbs (Figs 8A and 9); small tubercles present on temporal region, tympanum smooth, tympanic rim distinct but not elevated relative to skin of temporal region; on dorsal surfaces tubercles arranged in distinct longitudinal ridges on upper forelimbs, forearms, shanks and thighs, becoming less distinct on dorsal surfaces of hands, feet and digits (Fig. 9); six large tubercles on left flank and seven large tubercles on right flank irregularly scattered from axilla to groin, intermixed with smaller tubercles; central portion of outer edge of upper eyelid slightly thickened, with a distinct small single tubercle on a thickened ridge (Figs 8C, 9); distinct thick glandular supratympanic fold, narrow anteriorly, considerably widening posteriorly, extending from posterior corner of eye gently sloping down towards dorsal margin of tympanum (but not concealing it), where it broadly

curves down, terminating above axilla (Figs 8C, 9); short dorsolateral glandular ridge present above shoulders, on anterior part of dorsum, its length comparable with eye diameter (Fig. 9); a weak “>—<”-shaped glandular dermal parietoscapular-sacral ridge present on dorsum (Figs 8A, 9); transverse fold at head basis absent; dermal projection above cloaca absent; gular region, chest, abdomen and ventral surfaces of limbs smooth to weakly shagreened (especially on the posterior surface of abdomen); nuptial pad present, single, covered with microgranules, covering entire dorsal metacarpal of first finger extending distally to ca. 3/4 basal phalange length; pectoral glands round, flat, of medium size, positioned level with axilla; femoral gland flat, indistinct, on posterior surface of thighs.

Color of holotype in life. Entire dorsum light olive-brown to yellow-brown with large irregular brownish grey spots; dorsal surfaces of head yellowish brown from tip of snout to eyes; small oval-shaped spot with irregular borders on dorsal surface of snout between anterior canthi; a small dark dot on dorsal surface of snout tip; similar single dark dots on anterior parts of upper eyelids; brown “V”-shaped marking on crown between supraorbital horns with apex pointing posteriorly, outlined with thin light-beige edging; round brownish spot at head basis; “>—<”-shaped marking surrounded with dark olive-brown, outlined with thin light-beige edging forming a hourglass-shaped dorsal marking (Fig. 8A); two small roundish brown spots at sacrum (Fig. 8A); supratympanic fold dorsally light yellowish brown, ventrally dark-brown; front and lateral surfaces of snout and lateral canthus rostralis dark reddish brown; lateral surfaces of maxilla dark brown with four distinct orange-brown to yellowish beige bars extending from orbits towards edge of maxilla: smallest anteriormost light band borders nostril ventrally, with two posterior light bands extend from posterior corner of eye towards angle of mouth (Fig. 9); axilla purplish brown; tympanum uniform purplish brown; temporal region uniform dark purplish brown, clearly defined from light beige area on posterior part of maxilla; pupil black, outlined in copper-gold; iris golden dorsally and ventrally, copper-orange at medial part, with tiny dark reticulations spreading from pupil; sclera lemon-yellow; upper surface of limbs yellowish brown with irregular dark-brown spots on forearms and transverse spots forming dark-brown and greyish bands across shanks, thighs and tibio-tarsus (three complete transverse bands on left leg, two complete [both on shank and thigh] and one incomplete [on shank only] transverse bands on right leg: Figs 8 and 9), knee joint dark brown; sides beige-yellow with indistinct greyish white flecking and large black spots with irregular borders: marking location of large warts on each side of body, four smaller spots located dorsally, and two large brown-black spots located ventrally, top of larger flank tubercles brownish cream (Fig. 9); throat brownish to purplish grey with greyish white flecking and irregular dark-brown spots; chest and anterior half of abdomen purplish grey with whitish flecking and grey-brown blotches; posterior half of abdomen lighter greyish pink with irregular dark blotches; lower surface of limbs purplish grey with white and beige flecking; area surrounding vent and posterior surface of thighs dark black-brown with whitish spots, posterior surface of thighs near tibio-tarsal articulation black-brown with sparse whitish dots; dorsal surface of feet and shanks yellowish beige with brown flecking; ventral surface of feet and shanks brown-black. Pectoral and femoral glands creamy white. Nuptial pad, outer metacarpal (palmar) and metatarsal tubercles pink-

red to orange-red. Asperities covering dorsal surfaces of body, head, limbs and digits, lateral sides of head and anterior part of chest in life bright orange-red, forming reddish rows and ridges on dorsal surfaces of limbs as well as on edge of upper eyelid, palpebral projection also with orange-red asperities.

Color of holotype in preservative. In preservative coloration faded to light grey-brown on dorsum and flanks, with slightly paler limbs and greyish beige to whitish on venter; reddish and orange tints, as well as iris coloration, faded completely; dark markings on dorsum, sides and venter and other features remain without significant change (Fig. 8). Banding on limbs is less pronounced. Chest, abdomen, throat, interior portions of forelimbs and thighs are pale greyish brown; formerly brightly colored dorsal asperities and nuptial pads, palmar and metatarsal tubercles turned transparent or creamy white (Fig. 8C, E).

Measurements of the holotype (all in mm, taken by NAP). SVL 27.2; HW 8.0; HL 7.9; IFE 3.2; IBE 6.8; ED 3.6; TYD 1.9; TYE 1.6; SL 2.5; EN 1.3; NS 1.3; IUE 2.3; IN 2.3; UEW 2.0; FAL 7.0; HAL 7.3; FIL 2.4; FIIL 2.8; FIIL 4.6; FIVL 2.8; SHL 13.2; TL 12.1; FOL 11.9; TFOL 18.9; IMT 1.8; TIL 1.7; TIIL 3.8; TIIL 5.5; TIVL 6.7; TVL 3.5.

Variation. Morphometric variation within the type series and other referred specimens of the new species is shown in Table 5. Individuals of the type series are similar in morphology and body proportions (Figs 9, 10). There is a clear and significant difference in body size between males and females (Fig. 3): females (SVL 35.1–36.5 mm, $N = 6$) are significantly bigger than males (SVL 26.9–33.9 mm, $N = 29$) (Duncan test, $p < 0.05$); sexual differences were not significant for other mensural characters possibly due to the small sample size of females. Certain variation is observed in finger lengths: most of the examined specimens have the finger length formula $F1 < F4 < F2 < F3$ ($N = 14$), in some specimens, including the holotype, the second and the fourth fingers are of equal length ($F1 < F4 = F2 < F3$; $N = 5$), or the fourth finger is longer than the second ($F1 < F2 < F4 < F3$; $N = 5$); rarely the second finger is as long as the first finger ($F1 = F2 < F4 < F3$; $N = 2$). Specimens vary in the number and size of black spots and blotches on flanks (Fig. 10A, B). In life, both sexes of the new species have lighter dorsum and belly coloration when nocturnally active. Other in-life variation was observed for throat coloration: throat can be dark brownish with clear dark-grey blotches (Fig. 10C) to almost uniform brown-violet to purple with dark blotches not discernable (Fig. 10D). There is significant variation in dorsal pattern: in some specimens the dorsum looks almost uniform yellowish brown with an indistinct hourglass-shaped figure (Figs 6G, 10A, B, 14 [right]) whereas in other specimens the hourglass-shaped figure is distinct, dark brown and edged with light beige (Figs 6H, 9). There is some variation in the length of palpebral projections, from a small almost indistinct tubercle (Figs 6H, 13C) to a moderately well-developed projection (Figs 6G, 14). Coloration of the lateral surfaces of the head vary, but on all specimens two light suborbital bars are distinct, clearly separated from the uniform dark-brown coloration of the tympanal area. Iris coloration shows insignificant variation: Nui Chua Mt. population appear to have copper-red coloration of the entire iris (Fig. 14B), somewhat different from the coloration of the holotype (Fig. 9). Recently metamorphosed and

juvenile specimens have numerous bright red-orange tubercles (Fig. 10E, F) which are more conspicuous than in adults. Excluding the presence of nuptial pads on males, the new species shows no significant variation in dermal characters among sexes (Fig. 10); in preservative smaller tubercles become flattened and less distinct.

Tadpole description. Tadpoles were allocated to *Ophryophryne elfina* sp. n. based on the following evidence: (1) morphological features characteristic for megophryine larvae in general; *Ophryophryne* or *Megophrys* s. lato in particular (elliptical shaped body with long muscular tail, oral disk forms a dorsally oriented funnel); (2) collected in the stream where calling males of the new species were recorded; (3) species identification confirmed by mtDNA sequences of short 16S rRNA gene fragment (up to 500 bp) (GenBank Accession numbers: KY515232–KY515233, see Table 1).

Standard tadpoles measurements (mean \pm SD, $N = 5$, Stage 25; ZMMU A-5679, field number NAP-01169, collected from Bidoup Mt., 1900 m a.s.l., Bidoup–Nui Ba N.P., Lam Dong Prov.) (all in mm, taken by NAP): TOL = 28.4 ± 1.3 (27.4–30.2); BL = 8.6 ± 0.1 (8.4–8.7); TAL = 19.8 ± 1.2 (18.9–21.5); BW = 4.4 ± 0.4 (3.8–4.6); BH = 3.6 ± 0.2 (3.4–3.7); TH = 4.5 ± 0.4 (4.0–4.8); SVL = 9.2 ± 0.3 (9.0–9.5); SSp = 4.8 ± 0.2 (4.5–4.9); UF = 1.4 ± 0.1 (1.3–1.5); LF = 1.1 ± 0.0 (1.1–1.1); IN = 2.6 ± 0.1 (2.5–2.6); IP = 2.8 ± 0.2 (2.7–3.2); RN = 1.7 ± 0.1 (1.7–1.8); NP = 0.8 ± 0.1 (0.7–0.8); ED = 0.8 ± 0.0 (0.8–0.9).

The following description is based on five tadpoles at stage 25 (ZMMU A-5679, field number NAP-01169). In lateral view (Fig. 11A), body slightly compressed dorsoventrally (BH/BW $83.5 \pm 4.09\%$), especially anteriorly, convex both dorsally and ventrally. Body elliptical in dorsal view (Fig. 11B), with maximum width at middle of body (BW/BL $51.0 \pm 4.2\%$); snout short, rounded, blunt. Eyes of moderate size (ED/BL $9.8 \pm 0.3\%$), not bulging, separated by a distance which equals approximately 1.1 times internarial distance (IP/IN $110.2 \pm 6.8\%$), directed and positioned dorsolaterally, not visible in ventral view; pupils oriented dorsolaterally. Nares tubular, positioned dorsally (near anterior edge of eye), much closer to pupils than to tip of snout; directed laterally. Spiracle sinistral, conical, very short, opening at half of distance from snout tip to vent (SSp/SVL $52.2 \pm 3.4\%$); spiracle attached to body wall for most of its length, extremity is free, positioned at the level of longitudinal axis, oriented dorsoposteriorly, opening varies from rounded to oval. Tail long, more than two times longer than body (TAL/BL $231.3 \pm 11.0\%$), lanceolate; almost equal in height along its length (point of maximum height of tail located just anterior to midlength of tail); tail tip bluntly rounded, without terminal filament; tail musculature strong, gradually tapering, almost reaching tail tip. Tail fins shallow, moderately well developed, not extending onto body: dorsal fin originating almost at body-tail junction, much shorter than lower fin proximally and nearly equal in height to it on middle of tail; dorsal fin slightly higher than ventral fin on distal half of tail (LF/UF $77.1 \pm 4.9\%$); free margin of dorsal fin horizontal and shallow on anterior half of tail; free margin of ventral fin parallel to tail musculature. Vent opening medial, tubular, directed posteriorly, not linked to ventral tail fin. Neither skin glands nor neuromasts visible in preservative, but neuromasts of the lateral line system are distinct in life (Fig. 12A) forming two curved lines running from snout towards orbits and along orbital margins




Figure 10. Paratypes of *Ophryophryne elfina* sp. n. in life. **A–D** Bidoup Mt., Bidoup–Nui Ba N.P., Lam Dong Prov., 2000 m a.s.l.: **A** ZMMU A-4788 (field number NAP-01449), male, dorsolateral view **B** ZMMU A-4788 (field number NAP-01455), female, dorsolateral view **C** ZMMU A-4788 (field number NAP-01449), male, ventral view **D** ZMMU A-4788 (field number NAP-01455), female, ventral view **E–F** Chu Yang Sin Mt., Chu Yang Sin N.P., Dak Lak Prov., 1800 m a.s.l.: **E** ZMMU A-5691 (field number ABV-00580), metamorph, dorsolateral view **F** ZMMU A-5691 (field number ABV-00581), metamorph, dorsolateral view. Photos by N.A. Poyarkov.

ventrally. Subterminal oral disk with lips expanded vertically forming a dorsally oriented funnel (Fig. 11B); lateral corners of funnel distinct; upper lip notably smaller than lower; lips lack keratodonts, but bear short, low ridges, more densely arranged on upper than on lower labium, arranged in 18–24 (mean = 22) longitudinal rows and from 2–4 (mean = 3) transverse rows on upper labium to 4–6 (mean = 5) transverse rows on lower labium. Marginal papillae absent. Width of expanded funnel comprises over 75% of body length (Fig. 12A) and just 30% when folded in preservation (Fig. 12B).

Table 5. Measurements of the *Ophryophryne elfina* sp. n. specimens. For museum accession numbers relative to specimen collection numbers (NAP, ABV, HB, ROM, CYS) see Appendix 1. For locality details relative to population number see Table 1 and Fig. 1. For abbreviations see Material and methods. All measurements are given in mm. Table is continued on the next two pages.

Specimen ID	Population	Sex	Type status	FAL	HAL	FIL	FIIL	FIHL	FIVL	SHL	TL	FOL	TFOL	IMT	IMT
NAP-02658	10	m	holotype	27.2	8.0	7.9	3.6	1.9	1.6	2.5	1.3	1.3	2.3	2.3	2.0
NAP-01455	6	m	paratype	33.9	10.1	9.8	3.8	2.3	1.9	2.7	1.3	1.3	2.8	3.1	2.9
NAP-01449	6	m	paratype	31.0	9.3	9.8	3.8	2.2	1.6	2.5	1.3	1.2	3.1	2.9	3.5
NAP-01450	6	m	paratype	29.2	9.0	9.0	3.7	2.3	1.4	3.0	1.2	1.4	3.2	2.5	3.2
NAP-01460	6	m	paratype	29.6	8.7	8.4	3.4	2.7	1.1	2.1	1.1	1.0	2.5	1.5	2.6
ABV-00316	8	m		28.9	8.8	9.1	3.8	2.6	1.6	2.2	1.3	1.1	2.8	1.8	3.1
HB-36-1	8	m		27.5	8.6	9.2	4.1	2.4	1.3	2.6	1.4	1.4	2.6	2.1	2.6
ROM-36525	11	m		26.9	7.6	8.1	3.6	1.9	1.3	2.9	1.3	1.2	2.3	1.7	2.8
ROM-36523	11	m		27.8	8.2	8.4	4.0	2.2	1.8	2.8	0.9	1.5	2.1	1.6	2.9
ROM-36529	11	m		28.3	7.9	8.6	4.3	2.2	1.6	2.4	1.1	1.5	2.8	2.0	2.7
NAP-01757	6	m		32.0	9.2	9.2	4.0	2.3	1.8	2.6	1.3	1.3	2.3	2.3	2.5
NAP-01782	6	m		31.1	9.3	9.1	3.8	2.2	1.7	2.3	1.1	1.0	2.8	2.6	7.9
NAP-01758	7	m		32.1	9.2	9.3	3.4	2.4	1.7	2.4	0.8	1.3	3.1	2.0	2.4
NAP-01871	7	m		32.8	8.8	9.6	3.7	2.2	1.3	2.7	0.9	1.1	2.9	1.6	2.4
NAP-01783	7	m		31.3	9.0	9.2	4.0	2.2	1.6	2.8	1.2	1.1	3.1	2.1	2.8
ABV-00471	6	m	paratype	30.7	8.5	8.5	3.7	2.3	1.3	2.6	0.9	1.4	2.7	2.4	2.7
ABV-00454	6	m	paratype	31.4	8.8	9.3	3.8	2.1	1.5	2.0	1.0	1.1	2.5	1.9	2.9
ABV-00472	6	m	paratype	30.8	8.9	9.3	3.3	2.3	2.1	2.3	1.6	1.0	3.2	2.1	2.6
ROM-36524	11	m		28.5	8.3	8.5	3.6	2.1	1.2	2.8	0.8	1.6	2.4	2.2	2.5
ROM-36527	11	m		27.7	7.4	7.8	3.1	2.0	1.7	3.0	1.3	1.8	2.7	2.1	2.7
ROM-36522	11	m		28.1	7.9	8.4	3.2	1.7	1.3	2.8	1.2	1.5	2.6	2.1	2.8
ROM-36528	11	m		28.2	7.2	7.4	3.1	1.6	1.5	3.3	1.0	1.8	3.4	1.9	2.9
ROM-36526	11	m		27.1	7.6	8.2	3.6	2.1	1.5	2.9	1.1	2.1	2.6	2.1	2.4
ITBCZ 2828	8	m		27.9	8.2	8.9	3.9	2.2	1.2	2.5	1.1	1.4	2.1	2.8	3.1
ITBCZ 2908	8	m		29.4	9.6	9.1	3.4	2.3	1.4	3.3	1.1	2.2	2.7	3.3	2.7
ITBCZ 2909	8	m		30.1	9.5	9.8	4.1	2.2	1.8	2.9	1.3	1.6	2.4	3.3	2.7

Specimen ID	Population	Sex	Type status	FAL	HAL	FIL	FIIL	FIHIL	FIVL	SHL	TL	FOL	TFOL	IMT	IMT
ITBCZ 2918	8	m		29.6	8.7	9.3	3.4	2.2	1.3	2.8	1.1	1.7	2.2	2.9	2.7
ITBCZ 2919	8	m		30.8	9.2	9.6	3.4	2.1	1.6	3.3	1.4	1.9	2.4	2.6	2.6
ITBCZ 3502	8	m		28.6	8.7	8.9	3.5	2.3	1.8	3.3	1.2	2.1	2.6	2.8	2.5
ABV-00455	6	f	paratype	35.2	9.3	8.9	4.2	2.4	1.7	2.4	1.7	1.2	2.8	1.7	2.6
CYS-10-10	11	f		35.1	9.4	9.1	3.9	1.9	1.9	3.6	1.6	2.1	3.5	2.6	3.0
ROM-36530	11	f		35.7	9.0	8.6	3.6	2.1	1.6	3.3	1.0	1.9	3.6	2.4	2.8
ITBCZ 2786	8	f		36.5	10.0	11.0	4.1	2.4	1.8	2.7	1.2	1.5	2.7	2.8	3.1
ITBCZ 2788	8	f		35.8	10.1	10.9	4.2	3.0	1.6	3.2	1.3	1.9	2.3	2.6	2.8
ITBCZ 2792	8	f		35.3	10.2	10.3	4.1	2.2	1.8	3.3	1.5	1.8	2.8	2.9	3.1
Specimen ID	Population	Sex	Type status	FAL	HAL	FIL	FIIL	FIHIL	FIVL	SHL	TL	FOL	TFOL	IMT	IMT
NAP-02658	10	m	holotype	7.0	7.3	2.4	2.8	4.6	2.8	13.2	12.1	18.9	11.9	1.8	
NAP-01455	6	m	paratype	8.5	9.4	2.3	4.1	6.0	3.7	16.1	14.8	14.7	28.0	2.0	
NAP-01449	6	m	paratype	7.1	7.5	2.8	3.0	4.1	3.0	14.4	13.4	11.8	19.4	1.7	
NAP-01450	6	m	paratype	7.5	7.8	2.1	2.9	4.9	2.6	12.9	13.8	11.1	19.7	1.9	
NAP-01460	6	m	paratype	6.5	6.7	2.3	2.9	4.4	2.4	13.4	11.3	10.6	17.5	1.3	
ABV-00316	8	m		7.3	6.7	2.4	2.9	4.9	2.9	13.2	13.5	11.6	19.2	2.5	
HB-36-1	8	m		6.6	7.1	2.1	3.2	4.6	3.0	12.6	13.4	11.6	18.9	1.5	
ROM-36525	11	m		6.9	6.6	2.0	2.4	4.5	2.7	11.7	11.9	10.6	17.1	1.2	
ROM-36523	11	m		6.4	6.0	2.9	2.9	4.8	3.0	12.8	13.2	11.4	18.3	1.6	
ROM-36529	11	m		6.7	6.4	2.9	3.0	4.5	3.0	13.5	13.5	11.0	19.1	1.6	
NAP-01757	6	m		8.4	7.4	2.5	2.5	4.5	3.1	13.3	14.5	11.6	19.6	1.2	
NAP-01782	6	m		6.1	4.8	1.1	2.0	3.7	1.4	12.7	13.1	11.0	18.7	1.4	
NAP-01758	7	m		8.1	6.6	2.1	3.2	4.6	2.5	14.8	15.3	13.3	20.4	1.8	
NAP-01871	7	m		6.5	6.8	2.3	2.8	4.6	3.1	14.2	14.6	10.8	19.2	1.2	
NAP-01783	7	m		7.6	6.1	2.4	3.4	4.8	3.0	14.0	15.7	10.4	18.1	1.6	
ABV-00471	6	m	paratype	8.6	7.1	3.0	3.4	4.9	3.2	14.2	14.5	13.1	19.6	1.2	
ABV-00454	6	m	paratype	8.2	7.2	3.0	3.2	5.1	3.2	15.7	15.3	13.1	22.0	1.3	
ABV-00472	6	m	paratype	7.9	6.5	2.5	3.0	5.2	3.1	14.7	11.9	12.6	21.4	1.2	
ROM-36524	11	m		6.8	6.4	2.2	2.9	4.5	2.4	13.0	13.0	10.8	17.4	1.5	

ROM-36527	11	m		6.4	6.7	2.4	2.6	4.9	2.5	12.2	11.9	11.9	19.0	1.5
ROM-36522	11	m		6.8	6.6	2.2	2.7	4.3	2.3	12.3	12.9	11.3	18.7	1.2
ROM-36528	11	m		6.1	5.8	2.2	2.4	4.2	2.5	12.5	13.0	11.7	18.4	1.7
ROM-36526	11	m		6.1	6.3	2.3	2.5	4.3	2.7	12.7	12.4	12.7	18.2	2.2
ITBCZ 2828	8	m		7.8	7.1	3.2	2.9	4.1	3.4	4.6	12.0	11.2	19.6	1.8
ITBCZ 2908	8	m		7.5	8.0	3.5	2.8	5.1	3.9	14.8	14.9	12.7	22.8	1.9
ITBCZ 2909	8	m		7.8	8.2	3.1	3.8	5.2	4.1	16.3	16.1	12.6	21.9	2.1
ITBCZ 2918	8	m		7.0	7.3	3.4	3.3	5.0	3.5	19.6	13.4	12.1	20.0	1.7
ITBCZ 2919	8	m		8.9	8.3	3.4	3.0	6.0	3.6	16.0	15.1	12.3	21.9	2.1
ITBCZ 3502	8	m		7.8	7.7	3.2	3.0	5.5	3.5	15.0	14.7	12.6	20.7	1.8
ABV-00455	6	f	paratype	8.3	7.7	2.9	3.3	4.9	3.1	16.1	16.2	14.2	22.6	1.7
CYS-10-10	11	f		7.9	8.6	3.0	3.4	5.2	3.2	15.3	15.5	13.9	21.8	2.4
ROM-36530	11	f		7.7	8.7	3.3	4.7	5.6	3.6	14.9	15.8	15.1	23.3	1.5
ITBCZ 2786	8	f		9.8	9.2	4.1	3.9	6.3	4.6	19.8	17.8	15.7	27.2	2.3
ITBCZ 2788	8	f		10.1	9.7	4.2	4.1	7.0	4.9	17.7	15.5	15.3	25.0	2.0
ITBCZ 2792	8	f		10.1	9.1	4.5	4.1	5.8	5.2	19.0	17.2	15.1	26.0	2.1




Figure 11. External morphology of the tadpole of *Ophryophryne elfina* sp. n. (ZMMU A-5679, field number NAP-01169; Stage 25, TOL = 30.2 mm). **A** lateral view, mouth funnel closed (in preservative) **B** dorsal view, mouth funnel open (in life). Drawings by V.D. Kretova.

In life tadpoles have dorsal side of body and upper flanks uniform brownish red or brownish orange (Fig. 12A, B). Lower flanks weakly mottled with dark brown, few round blackish spots on tail and dorsum; with orange neuromasts visible on dorsal surface (Fig. 12A). Abdomen light brownish orange, intestine not visible through body wall. Caudal muscles pale; tail fins translucent with a few darker spots (more on upper than on lower fin); dorsally tail with indistinct middorsal orange line (Fig. 12A). Eyes golden with black reticulations. Oral funnel pinkish orange with brownish red papillae (Fig. 12B). In preservative tadpole coloration gets much duller, but the general coloration pattern is still visible after 7 years in ethanol.

Advertisement call characteristics. Refer to the Acoustic differentiation section, Table 4 and Fig. 5, for bioacoustic comparison of the new species with *O. synoria* and *O. gerti*. Refer to Appendix 3 for call variation data in *Ophryophryne elfina* sp. n.

Position in mtDNA phylogeny and sequence divergence. The new species is reconstructed as a member of the *Ophryophryne* Group II (Fig. 2), forming a sister group with respect to the clade joining *O. gerti* and *O. synoria* (see Fig. 2). Uncorrected genetic *p*-distances between *Ophryophryne elfina* sp. n. partial sequences for the 16S rRNA gene and all homologous sequences available on GenBank included in the analysis (see Table 1) varied from 8.2% (with *O. gerti* s. stricto, clade A) to 10.0% (with *O. hansii* and *O. synoria*, clade C) (see Table 2). This degree of pairwise divergence in the 16S rRNA gene is greater than that usually representing differentiation at the species level in anura (Vences et al. 2005a, 2005b, Vieites et al. 2009, Poyarkov et al. 2015a, 2015b). Intraspecific variation in this gene fragment for *Ophryophryne elfina* sp. n. is significant — maximum sequence divergence between Nui Chua Mt. population



Figure 12. Coloration of *Ophryophryne* tadpoles in life: **A** *Ophryophryne elfina* sp. n. “(ZMMU A-5684, field number NAP-02673; Stage 25) from Chu Pan Fan Mt., 1900 m a.s.l. in Chu Yang Sin N.P., Dak Lak Prov., mouth funnel open **B** tadpole from the same locality and collection data (ZMMU A-5684, field number NAP-02673; Stage 25) with mouth funnel closed.. Photos by N.A. Poyarkov.

(Fig. 2, clade D) and populations from the rest of the species range (Fig. 2, clade E) is $p = 3.1\%$. Intraspecific variation in 16S rRNA gene fragment within one geographic population was lower, and ranged from 0.0% to 0.8% of substitutions.

Distribution. *Ophryophryne elfina* sp. n. is found to be endemic to five provinces in (Lam Dong, Dak Lak, Khanh Hoa, Ninh Thuan and Phu Yen) in the northern and eastern part of the Langbian Plateau and its foothills in southern Vietnam (localities 6–12, Fig. 1). The new species is restricted to wet evergreen montane tropical and elfin forests, receiving high precipitation from the sea. Such wet forests are found only on high elevations in the central parts of the Langbian Plateau (e.g. 1900–2100 m a.s.l. on Bidoup Mt., Lam Dong Prov., Fig. 1, Loc. 6) or peripheral mountains remote from the sea (e.g. 1900–2300 m a.s.l. on Chu Pan Fan and Chu Yang Sin Mts., Dak Lak Prov., Fig. 1, Locs 10 and 11), but on the eastern foothills of the plateau which receive more precipitation, the new species is found at lower elevation (from 950 to 1510 m a.s.l. on Hon Ba Mt., Khanh Hoa Prov., Fig. 1, Loc. 8; 780 m a.s.l. on Nui Chua Mt., Ninh Thuan Prov., Fig. 1, Loc. 9; and 700 m in Phu Yen Prov., Fig. 1, Loc. 12).

Ecology. All specimens were collected at night after heavy rains along montane cascade rocky streams, along small waterfalls, or intermittent rocky brooks; or found during the day time under tree-logs and within leaf litter in the limited fragments of

primary montane wet polydominant evergreen tropical forests, with a high abundance of large rocks and fallen trees covered with a thick layer of mosses. This including high montane forests that are composed of the specific floral community known as “elfin” forests, with miniature trees (up to 10 m tall). These areas always have high precipitation and have much milder climate than other tropical forests in southern Vietnam: active breeding of the new species was recorded in February with temperatures of ca. 11–12°C.

On Bidoup Mt. summit (Lam Dong Prov.), *Ophryophryne elfina* sp. n. was recorded from 1890 to 2035 m a.s.l. in montane polydominant high canopy (trees up to 35 m tall) and elfin (trees up to 10 m tall) (sub)tropical forests with the predominance of trees of the family Fagaceae (*Lithocarpus* sp., *Castanopsis* sp.), Elaeocarpaceae (*Elaeocarpus* sp.), Lauraceae (*Machilus* sp.), Magnoliaceae (*Magnolia* sp., *Michelia* sp.), and occasional large trees of *Fokienia hodginsii* (Cupressaceae). These forests have thick leaf litter, numerous fallen logs and rocks covered with mosses, and an undergrowth that is predominated by ferns (mostly *Asplenium* sp., Aspleniaceae) (see Kuznetsov and Kuznetsova 2011) (Fig. 15A). On Hon Giao mountain ridge, the new species was found along mountain streams from 1800 to 1900 m a.s.l. in montane polydominant elfin forests with the predominance of trees of the family Fagaceae (*Castanopsis* sp., *Lithocarpus* sp.), Lauraceae (*Cinnamomum* sp., *Neolitsea* sp.), Ericaceae (*Rhododendron* sp.), Magnoliaceae (*Mangletia* sp., *Michelia* sp.), Elaeocarpaceae (*Elaeocarpus* sp.) and Podocarpaceae (*Podocarpus neriifolius*). This forest had a dense undergrowth of mosses, orchids (*Coelogyne* sp., *Dendrobium* sp.; Orchidaceae) and occasional ferns (*Cyathea* sp., Cyatheaceae) (see Kuznetsov and Kuznetsova 2011) (Fig. 15B). In Chu Yang Sin N.P. (Dak Lak Prov.), the new species was found from 1800 to 2100 m a.s.l. in montane forests with the predominance of trees of the families Pinaceae (*Pinus krempfii*; *Pinus kesiya*) and Fagaceae (*Lithocarpus* sp., *Castanopsis* sp.), with dense undergrowth of ferns, numerous rocks and fallen trees covered with mosses (Fig. 15C). In Hon Ba N.R. (Khanh Hoa Prov.), the new species was found from 950 to 1510 m a.s.l. along mountain streams in forests having polydominant composition including Fagaceae (*Lithocarpus* sp., *Quercus* sp.), Elaeocarpaceae (*Elaeocarpus* sp.), Theaceae (*Thea* sp., *Camellia* sp.), Lauraceae (*Cinnamomum* sp., *Neolitsea* sp.), Araliaceae (*Schefflera* sp.) and Rutaceae (*Euodia* sp.).

On Bidoup Mt. summit (1890–2035 m a.s.l.; Lam Dong Prov.) *Ophryophryne elfina* sp. n. occurs in syntopy with *Leptobrachium pullum* (Smith, 1921), *Leptobrachium leucops* Stuart, Rowley, Tran, Le & Hoang, 2011, *Leptolalax bidoupensis* Stuart, Rowley, Tran, Le & Hoang, 2011, *Leptolalax pallidus* Rowley, Tran, Le, Dau, Peloso, Nguyen, Hoang, Nguyen & Ziegler, 2016, *Ingerophrynus galeatus* (Günther, 1864), *Hylarana montivaga* (Smith, 1921), *Rhacophorus vampyrus* Rowley, Le, Thi, Stuart & Hoang, 2010, *Theloderma palliatum* Rowley, Le, Hoang, Dau & Cao, 2011 and *Raorchestes gryllus* (Smith, 1924). On Hon Giao Mt. (1900–2000 m a.s.l.; borders of Lam Dong and Khanh Hoa provinces), *Ophryophryne elfina* sp. n. occurs in syntopy with *Leptobrachium leucops*, *Leptolalax bidoupensis*, *Duttaphrynus melanostictus* (Schneider, 1799), *Hylarana montivaga*, *Rhacophorus vampyrus* and *Raorchestes gryllus*. On Chu Pan Fan and Chu Yang Sin Mts. (1900 m a.s.l., Dak Lak Prov.), the new species is found in syntopy with *Xenophrys* cf. *maosonensis* (Bourret, 1937), *Leptobrachium*




Figure 13. Comparison of the head coloration in life of three *Ophryophryne* species from the Langbian Plateau: **A** *Ophryophryne synoria*, Bu Gia Map N.P., Binh Phuoc Prov., Vietnam **B** *Ophryophryne gerti*, Chu Yang Sin N.P., Dak Lak Prov., Vietnam **C** *Ophryophryne elfina* sp. n., Hon Giao Mt., Bidoup–Nui Ba N.P., Lam Dong Prov. Photos by N.A. Poyarkov and N.L. Orlov.

sp., *Leptotalax* sp., *Hylarana montivaga*, *Rhacophorus vampyrus* and *Raorchestes gryllus*. On Chu Yang Sin Mt. (1700–1800 m a.s.l., Dak Lak Prov.), the new species is also found in syntopy with *Ophryophryne gerti* (Fig. 14A). On Hon Ba Mt. (1500 m a.s.l., Khanh Hoa Prov.), the new species was recorded in syntopy with *Leptobrachium leucops*, *Leptotalax* sp., *Microhyla arboricola* Poyarkov, Vassilieva, Orlov, Galoyan, Tran, Le, Kretova & Geissler, 2014, *Theloderma truongsongense* (Orlov & Ho, 2005) and *Raorchestes gryllus*.

Reproductively active males were found while calling along streams, usually sitting on leaves of ferns or on the stone banks, rarely on rocks or large stones (see Fig. 14B, C). Some specimens were collected hiding amongst fern stems and were difficult to locate. Females were found hiding under tree logs or in the forest litter.

The ovaries of females contained well-developed unpigmented eggs with a diameter of approximately 2.2–2.8 mm ($N = 15$; measured from ZMMU ABV-00455, gravid female). On Hon Ba Mt., calling males were observed between 22 to 24 December, 22 to 28 March and 15 to 18 October. On Bidoup, Hon Giao and Chu Yang Sin Mts., reproductive activity and calling males were recorded from 10 February until mid-July. Tadpoles were found from April until July in the same streams where calling males were recorded; during the day time tadpoles hide under flat stones or dead leaves on the stream bed, but come out and can be visible in the shallow sandy parts of the stream at night. Metamorphosed individuals were observed in Chu Yang Sin N.P. in May.

Conservation status. The full extent of the distribution of *Ophryophryne elfina* sp. n. is unknown, and the discovery of new localities on mountain ridges at elevations above 1500 m are highly anticipated. To date the species' range includes the following nature conservation areas: Bidoup–Nui Ba N.P. (Lam Dong Prov.), Chu Yang Sin N.P. (Lam Dong Prov.), Hon Ba N.R. (Khanh Hoa Prov.) and Nui Chua N.P. (Ninh Thuan Prov.). However, population size and population dynamics of the new species are unknown. Given the available information, we suggest *Ophryophryne elfina* sp. n. to be considered as a Data Deficient species following IUCN's Red List categories (IUCN Standards and Petitions Subcommittee 2016).

Comparisons. *Ophryophryne elfina* sp. n. is one of the smallest species of its genus, with adult male size (SVL 26.9–33.9 mm) similar to that of *O. pachyproctus* (adult male SVL 28.0–30.0 mm).

Ophryophryne elfina sp. n. differs from allopatric *O. hansi* (central Vietnam and neighboring southeastern Laos and northeastern Cambodia; Ohler 2003, Stuart 2005, Stuart et al. 2006) by its much smaller adult size: *Ophryophryne elfina* sp. n. male SVL 26.9–33.9 mm, $N = 29$, female SVL 35.1–36.5 mm, $N = 6$ (vs. *O. hansi* male SVL 33.4–43.1 mm, $N = 12$, female SVL 45.1–53.9 mm, $N = 5$; our data), by skin of dorsal and lateral surfaces of head, body and limbs shagreened with numerous small tubercles (vs. notably tubercular and warty skin on dorsal surfaces), and short dorsolateral glandular ridge above each shoulder (vs. dorsolateral glandular ridges absent).

Ophryophryne elfina sp. n. differs from allopatric *O. pachyproctus* (Yunnan Prov. in southern China, central Vietnam and possibly adjacent regions of Laos; Bain et al. 2007, Fei et al. 2009, 2010, 2012) by lacking dermal protuberance with dermal flaps




Figure 14. *Ophryophryne elfina* sp. n. in situ: **A** Two syntopically collected males of *Ophryophryne gerti* (left) and *Ophryophryne elfina* sp. n. (right) in Chu Yang Sin N.P., Dak Lak Prov., Vietnam, 1750 m a.s.l., photo by N.L. Orlov **B** calling adult male of *Ophryophryne elfina* sp. n. in Nui Chua Mt., Nui Chua N.P., Ninh Thuan Prov., Vietnam, 780 m a.s.l., photo by S.N. Nguyen **C** adult male of *Ophryophryne elfina* sp. n. in calling position in Hon Ba N.R., Khanh Hoa Prov., Vietnam, 1510 m a.s.l., photo by L.T. Nguyen.

above cloacal opening (vs. present on *O. pachyproctus*), short dorsolateral glandular ridge above each shoulder not connected to posterior tips of “><”-shaped glandular parietoscapular-sacral ridge (vs. dorsolateral glandular ridge connected to posterior tips of “H”- or “Y”-shaped glandular parietoscapular-sacral ridge), supratympanic fold light brown dorsally on males (vs. white or light beige dorsally on males), and males with nuptial pad on first finger only (vs. one nuptial pad each on first and second fingers).



Figure 15. Natural habitat of *Ophryophryne elfina* sp. n. on Langbian Plateau, southern Vietnam. **A** Elfin forest on the top of Bidoup Mountain (ca. 2100 m a.s.l.), Bidoup–Nui Ba N.P., Lam Dong Prov. **B** elfin forest on the top of Hon Giao Mountain (ca. 2000 m a.s.l.), Bidoup–Nui Ba N.P., border of Lam Dong and Khanh Hoa provinces **C** typical breeding site along a mountain stream in an evergreen mixed sub-tropical montane forest on northern slopes of the Chu Pan Fan Mountain, Chu Yang Sin N.P., Dak Lak Prov. (type locality) (ca. 1900 m a.s.l.). Photos by N.A. Poyarkov and O.V. Morozova.

Ophryophryne elfina sp. n. differs from allopatric *O. microstoma* (Guangxi and Yunnan Provs., southern China to northern Vietnam, and northern Thailand; Khonsue and Thirakhupt 2001, Chan-ard 2003, Fei et al. 2009, 2010, 2012, Nguyen et al. 2009), by much smaller adult size: *Ophryophryne elfina* sp. n. male SVL 26.9–33.9 mm, $N = 29$, female SVL 35.1–36.5 mm, $N = 6$ (vs. *O. microstoma* male SVL 34.3–44.4 mm, $N = 10$, female SVL 39.4–57.0 mm, $N = 7$; Ohler 2003, Fei et al. 2009, Stuart et al. 2010; our data), short anterior dorsolateral glandular ridge above each shoulder (vs.

dorsolateral ridges long, ca. 65–90% trunk length), and large tubercles posteriorly on dorsal surface of body (vs. large tubercles absent).

Ophryophryne elfina sp. n. differs from allopatric *O. poilani* (found in the mountains of the Tay Nguyen Plateau of central Vietnam and in adjacent areas of northeastern Cambodia, and, possibly, Laos; Bourret 1937, Stuart et al. 2010) by having smaller adult body size: *Ophryophryne elfina* sp. n. male SVL 26.9–33.9 mm, $N = 29$, female SVL 35.1–36.5 mm, $N = 6$ (vs. *O. poilani* male SVL 32.6–38.1 mm, $N = 14$, female SVL 47.4–50.8 mm, $N = 2$; Stuart et al. 2010), short dorsolateral glandular ridge above the shoulder not connected to posterior tips of “>—<”-shaped parietoscapular-sacral glandular ridge (vs. dorsolateral glandular ridge connected to posterior tips of “>—<”-shaped parietoscapular-sacral glandular ridge), and lacking characteristic dark “mask-like” coloration of temporal region, and supratympanic fold being dark brown ventrally and light brown dorsally on males (vs. temporal region and lateral surfaces of head uniformly dark-colored forming a dark “mask”, extending posteriorly towards axilla; supratympanic fold edged with white on males).

Ophryophryne elfina sp. n. differs from sympatric *O. synoria* (found at lower elevation from 200 to 1500 m a.s.l. in the foothills of the Langbian Plateau in southern Vietnam and adjacent easternmost hilly Cambodia) by much smaller adults body size: *Ophryophryne elfina* sp. n. male SVL 26.9–33.9 mm, $N = 29$, female SVL 35.1–36.5 mm, $N = 6$ (vs. *O. synoria* male SVL 38.2–53.7 mm, $N = 14$, female SVL 51.4–70.7 mm, $N = 3$; our data; Fig. 3), red-orange nuptial pad (in life) on first finger only (vs. two nuptial pads, covered in brown microgranules, large on first finger, covering entire dorsal metacarpal extending to 3/4 basal phalange length, on second finger medium sized on metacarpal extending to mid basal phalange on inner dorsal side), numerous bright red-orange asperities (in life) on dorsal and lateral surfaces of body, head and dorsal surfaces of limbs (vs. black and white asperities, small sized, spinular, moderately dense in narrow band along lower jaw, and on posterior upper jaw, few on tympanic region [excluding tympanum], along supratemporal folds and on posterior upper eyelids; some on anterior dorsum, becoming moderately dense posteriorly, above and surrounding cloaca, few on dorsal shanks, and absent on remaining surfaces on holotype of *O. synoria*), and smaller tympanum/eye diameter ratio, TYD/ED 48.9%–62.6%, $N=29$ (vs. TYD/ED 64.8%–85.2%, $N=14$).

Ophryophryne elfina sp. n. differs from sympatric *O. gerti* (found at mid-elevations from 700 to 2000 m a.s.l. in the central and northern parts of the Langbian Plateau in southern Vietnam) by typically smaller adults body size: *Ophryophryne elfina* sp. n. male SVL 26.9–33.9 mm, $N = 29$, female SVL 35.1–36.5 mm, $N = 6$ (vs. *O. gerti* male SVL 31.7–42.2 mm, $N = 15$, female SVL 43.1–47.4 mm, $N = 3$; our data; Fig. 3), bright red-orange nuptial pads on males in life (vs. grey or black-brown nuptial pads on males in life), short dorsolateral glandular ridge above each shoulder, not connected to posterior tips of “>—<”-shaped parietoscapular-sacral glandular ridge, see Figs 6G–H, and 8 (vs. strong dorsolateral glandular ridge from above each shoulder to approximately 4/5 distance between axilla and groin, connecting with posterior tips of “>—<”-shaped parietoscapular-sacral glandular ridge; see Figs 6A–C, 7), skin on dor-

sal and lateral surfaces of body shagreened with numerous small tubercles (vs. skin on dorsum and sides of body granular, with numerous small and medium-sized tubercles and larger warts, see Fig. 6A–C), dark hourglass-shaped markings on dorsum normally edged with white, see Figs 6H and 8 (vs. dark hourglass-shape on dorsum indistinct or, if present, unclear and not edged with white, Fig. 6A–C), and throat, chest and abdomen having generally lighter coloration than in *O. gerti*.

Though available information on tadpole morphology of *Ophryophryne* is very limited (Liu and Hu 1962, Huang et al. 1991, Grosjean 2003, Fei et al. 2009), the tadpoles assigned to the new species based on the analysis of short 16S rRNA gene sequences (Table 1) have certain morphological characteristics that could be useful for distinguishing the larval stage of *Ophryophryne elfina* sp. n. from other *Ophryophryne* species. From tadpoles of *O. microstoma*, described in detail by Grosjean (2003), tadpoles of *Ophryophryne elfina* sp. n. differ mainly by possessing a longer tail: TOL/BL ratio $231.3 \pm 11\%$, $N = 5$ (vs. TOL/BL $< 210\%$, $N = 52$ on *O. microstoma*), and tail tip rounded (vs. tail tip bluntly pointed), mean = 22 longitudinal rows of papillae and from 2–4 (mean = 3) transverse rows of papillae on the upper labium and 4–6 (mean = 5) transverse rows of papillae on the lower labium, $N = 5$ (vs. mean = 20 longitudinal rows of papillae, and two upper labium and four lower labium transverse rows of papillae [without clear limits], $N = 52$); however, some of these differences may relate to the fact that Grosjeans' description was based on later developmental stages (Gosner stage 37) than our sampling (Gosners' stage 25).

DNA-barcoding using short sequences for 16S rRNA (Table 1) also enabled us to identify tadpoles of sympatric *Ophryophryne* species from the Langbian Plateau, and though our sampling is not big enough to provide detailed morphological descriptions of larval morphology for *O. gerti* and *O. synoria*, we found some differences in coloration of tadpoles which may be useful for preliminary diagnostics of the three sympatric *Ophryophryne* species in the wild. Despite overall morphological similarity, both *O. gerti* and *O. synoria* show the presence of light golden to copper blotches on dorsal surfaces of the body and tail, whereas *Ophryophryne elfina* sp. n. tadpoles always have distinctive uniform brownish coloration with small coppery dots (Fig. 12A–B).

Despite overall similarity, advertisement calls of each *Ophryophryne* species inhabiting the Langbian Plateau are easily diagnosable based on acoustic parameters. Some parameter values clearly differ between all of the studied species (Fig. 5). For example, the highest values of the repetition rate per recording and per series are found for *O. synoria* (3.07 ± 0.13 calls/s, $N = 3$, and 5.34 ± 0.15 calls/s, $N = 15$, respectively) while the lowest are reported for *O. gerti* (0.35 ± 0.14 calls/s, $N = 3$, and 2.33 ± 0.03 calls/s, $N = 108$, respectively). *Ophryophryne elfina* sp. n. occupies an intermediate position between these two species (1.18 ± 0.2 calls/s, $N = 5$, and 3.87 ± 0.07 calls/s, $N = 140$, respectively; differences significant, $F_{2,8} = 46.7$, $p < 0.001$ and $F_{2,260} = 220.7$, $p < 0.001$, respectively, one-way ANOVA).

The call temporal parameters for *Ophryophryne elfina* sp. n. compared to sympatric *O. synoria* and *O. gerti*, also differ for series duration, which is the highest in calls of *Ophryophryne elfina* sp. n., comprising 3.42 ± 0.11 s, $N = 140$ (see Table 4) (differences

with *O. gerti* significant, $F_{2,267} = 40.4$, $p < 0.001$, one-way ANOVA; differences with *O. synoria* not significant, see Table 4 for details). The advertisement call of the new species is further significantly different from calls of *O. gerti* for values of a number of acoustic parameters (see Table 4), such as the number of calls per series (12.84 ± 0.41 , $N = 140$, versus 4.64 ± 0.16 , $N = 115$, for *O. gerti*; differences significant, $F_{2,267} = 151.4$, $p < 0.001$, one-way ANOVA), the call duration (73 ± 0.23 ms, $N = 1797$, versus 104 ± 0.5 ms, $N = 533$, for *O. gerti*; differences significant, $H_{2,2530} = 1345.1$, $p < 0.001$, Kruskal-Wallis ANOVA) and inter-calls interval (207 ± 2.06 ms, $N = 1657$, versus 421.54 ± 4.17 ms, $N = 418$, for *O. gerti*; differences significant, $H_{2,2260} = 1008.5$, $p < 0.001$, Kruskal-Wallis ANOVA). The advertisement call of *Ophryophryne elfina* sp. n. is further different from calls of *O. synoria* in the frequency of maximum amplitude (4645.94 ± 4.39 Hz, $N = 1797$, versus 3798.9 ± 4.87 Hz, $N = 200$; differences significant, $H_{2,2530} = 1030.2$, $p < 0.001$ ($U = 0$), Kruskal-Wallis ANOVA).

Finally, the new species is markedly distinct from all other congeners for which comparable sequences are available, including its closest relatives *O. gerti* and *O. synoria*, by relatively large genetic distances in 16S rRNA mtDNA gene fragment ($p \geq 8.2\%$).

Discussion

The data presented here provide the most extensive molecular sampling for the elucidation of phylogenetic relationships within the genus *Ophryophryne*. According to our data, genetic variation within *Ophryophryne* appears to be strongly geographically structured. Thus, our results indicate the division of the genus *Ophryophryne* into two major reciprocally monophyletic groups: one corresponding to species found on the Langbian Plateau (Group II, Fig. 2), and another joining species found outside the plateau from central and northern Truong Son and adjacent areas (Group I, Fig. 2). Our data support the hypothesis that eastern Indochina, including the central and southern parts of the Truong Son Mountains (known also as Tay Nguyen Plateau), host the highest diversity of *Ophryophryne*, and was the center of radiation for this genus (Orlov and Ananjeva 2007, Mahony et al. 2017). Similar patterns of geographic structuring of mtDNA lineages were reported for the genus *Leptolalax*, another megophryid genus inhabiting the Truong Son Mountains (Poyarkov et al. 2015a, Rowley et al. 2016).

A hidden diversity of *Ophryophryne* frogs is revealed in the mountains of the Langbian Plateau, where previously only one species, *O. gerti*, was correctly reported (Ohler 2003, Nguyen et al. 2009, Stuart et al. 2010). In our study it is shown that the previous records of *O. cf. gerti* from central Vietnam and Laos (Ohler 2003, Bain et al. 2007) actually belong to different species of *Ophryophryne* and thus we clarify the range of *O. gerti* showing that this species is likely endemic to the Langbian Plateau. The known distribution of *O. synoria* is also extended, previously known exclusively from Cambodia (Stuart et al. 2006) and adjacent provinces of Vietnam (Vassilieva et al. 2016), and demonstrate that this species has a considerably wider range encompassing the central, northern and western edges of the Langbian Plateau. Finally, we describe the new spe-

cies *Ophryophryne elfina* sp. n., which is endemic to the northern and eastern edges of the plateau. Thus, the Langbian Plateau was a center of *Ophryophryne* radiation and cradles three endemic species of these frogs; all of them are sympatric in eastern and northern parts of the plateau and often can be recorded in synbiotopy.

Ophryophryne elfina sp. n. represents one of the smallest known species of the genus *Ophryophryne*. We found that the three *Ophryophryne* species of the Langbian Plateau are differentiated in body size with the largest species *O. synoria* preferring lowland and foothill monsoon forests at elevations from 200 to 1500 m a.s.l., medium-sized *O. gerti* found in evergreen montane tropical forests at mid-elevations from 700 to 2000 m a.s.l. and the smallest species *Ophryophryne elfina* sp. n. being restricted to wet montane subtropical forests at elevations from 700 to 2100 m a.s.l., including elfin forests at high elevations. It is probably not surprising that advertisement calls of the three occasionally sympatric *Ophryophryne* species show significant differences both in call structure and frequency parameters (see Table 4, Fig. 5), and the three studied species are characterized by relatively high values of the frequency parameters (as compared to several other Megophryidae species studied acoustically, especially of the genera *Leptolalax*, see review in Rowley et al. 2016, and *Leptobrachium*, see e.g. Stuart et al. 2010). The high frequency parameters may be related with their tendency to vocalize in close proximity to mountain cascade streams, which would create a low-frequency background noise (Preininger et al. 2007). It was shown that low background noise may induce frogs to call at higher frequency rates than expected from their body size, thereby improving the signal-to-noise ratio of their calls (Penna et al. 2005, Wells 2007, Goutte et al. 2016). Our study also recorded that the values of some temporal call parameters of *Ophryophryne elfina* sp. n. significantly differ between February (average temperature 11.3°C) and April recordings (average temperature 17.5°C; see Appendix 3 for details). Our results correspond with previous reports that intraspecific variation of temporal parameters of anuran calls can depend upon temperature (e.g., Gerhardt and Huber 2002).

The frequency of maximum amplitude coincides with the fundamental frequency for all *Ophryophryne* species, and have almost equal values for *Ophryophryne elfina* sp. n. and *O. gerti* (4645.94 ± 4.39 Hz, $N = 1797$, and 4845.99 ± 4.22 Hz, $N = 533$, respectively). The frequency of maximum amplitude of *O. synoria* is significantly lower (3798.9 ± 4.87 Hz, $N = 200$; see Table 4 for details), which may be related to the larger body size of the latter species (Stuart et al. 2006, Wells 2007). Further studies on acoustic communication of Langbian *Ophryophryne* species in areas of allopatry and sympatry would be valuable for better understanding the bioacoustic patterns observed here.

The Langbian Plateau is known for its high herpetofaunal diversity and endemism, a significant portion of which has been discovered only recently (e.g., Orlov et al. 2008, 2012, Rowley et al. 2010c, 2011a, 2011b, 2016, Stuart et al. 2011, Poyarkov [Paiarkov] and Vassilieva 2011, Nazarov et al. 2012, Chan et al. 2013, Hartmann et al. 2013, Geissler et al. 2014, 2015, Vassilieva et al. 2014, Poyarkov et al. 2014, 2015a, 2015b). Despite this increase in species discoveries, many areas of the Annamites have received little scientific attention and are very likely to host further

previously unknown diversity. The need for biological exploration in this region is made more urgent given the ongoing loss of natural habitats due to logging, road construction, increasing agricultural pressure and other human activities (Meijer 1973, De Koninck 1999, Laurance 2007, Meyfroidt and Lambin 2008, Kuznetsov and Kuznetsova 2011).

Habitat loss is the greatest threat to amphibians in southeast Asia, and the amphibians of the region appear to be particularly vulnerable to habitat alterations (Rowley et al. 2010b). Frogs of the genus *Ophryophryne* depend on fast-flowing clean mountain streams for reproduction, and appear to be restricted to relatively undisturbed broad-leaf evergreen forests: such habitat specialist range-restricted species are likely to be most at risk (Poyarkov et al. 2012, Rowley et al. 2010b, 2016). Deforestation, habitat loss and modification are continued threats in southern Indochina (Meyfroidt and Lambin 2008), and further studies on herpetofaunal biodiversity in this region are urgently required for elaboration of effective conservation measures.

Addendum (added post manuscript acceptance)

Due to the simultaneous review period of the present paper, and the now recently published Mahony et al. (2017), we chose to preliminarily use *Ophryophryne* at the genus level (following Chen et al. 2017; published online 1 December 2016), pending the publication of the taxonomic justification by Mahony et al. (2017) which supports a subgenus level classification of *Ophryophryne* within *Megophrys*. Mahony et al. (2017) also provided the replacement name *Megophrys (Ophryophryne) koui* Mahony, Foley, Biju & Teeling, 2017 for *Ophryophryne pachyproctus* Kou, 1985. We suggest that the new species combination *Ophryophryne elfina* sp. n. should hereafter be referred to as *Megophrys (Ophryophryne) elfina* (Poyarkov, Duong, Orlov, Gogoleva, Vassilieva, Nguyen, Nguyen, Nguyen, Che & Mahony) to reflect this revised taxonomy.

Author contributions

NA Poyarkov envisioned the original idea of the manuscript, collected material and data in the field and in the lab, executed this study and wrote the manuscript; TV Duong performed morphometric, molecular and phylogenetic analyses; NL Orlov collected material in the field; SS Gogoleva collected data in the field and performed acoustic analyses and wrote the relevant parts of the manuscript; AB Vassilieva collected material and data in the field; LT Nguyen collected material in the field and assisted with morphological analysis; VDH Nguyen, J Che and SN Nguyen collected material in the field and provided additional molecular data; S Mahony examined type and comparative specimens, performed molecular analysis, provided redescription of types, and edited the manuscript. All authors contributed to this paper sufficiently.

Acknowledgements

Fieldwork was funded by the Joint Russian-Vietnamese Tropical and Technological Center (JRV TTC) and was conducted under permission of the Bureau of Forestry, Ministry of Agriculture and Rural Development of Vietnam (permits Nos. 170/ TCLN–BTTN of 07/02/2013; 400/TCLN–BTTN of 26/03/2014; 831/ TCLN–BTTN of 05/07/2013) and of local administration (Lam Dong Prov.: No. 5832/UBND–LN of 22/10/12; Khanh Hoa Prov.: No. 522/SngV–TTDN&HTQT of 13/07/2013; fieldwork was conducted in accordance to the Agreement No. 37/HD on the scientific cooperation between Cat Tien N.P. and the JRV TTC and the Agreement № 137/HD NCKH of 23.06.2010 on the scientific cooperation between Bu Gia Map N.P. and the JRV TTC). The authors are grateful to Andrei N. Kuznetsov, Leonid P. Korzoun and Vitaly L. Trounov for support and organization of fieldwork. We sincerely thank our Vietnamese colleagues Nguyen Dang Hoi, Hoang Minh Duc, Nguyen Ngoc Hung, Nguyen Thien Tao, Pham Thi Ha Giang, Tran Tien and Le Xuan Son for help and continued support. Many thanks to Eduard A. Galoyan, Igor V. Palko, Evgeniy S. Popov, Olga V. Morozova, Alina V. Alexandrova, Valentina D. Kretova and Evgeniya N. Solovyeva for their help during the fieldwork, assistance in the laboratory and continued support of this project. We thank Ilya A. Volodin for help with acoustic analyses. For permission to study specimens under their care, we thank Valentina F. Orlova (ZMMU) and Natalia B. Ananjeva (ZISP). We are grateful to Andrey N. Kuznetsov for help with plant identification. We are sincerely grateful to Angelica Crottini, Annemarie Ohler and an anonymous reviewer for their kind help and useful comments which helped us to improve the previous version of the manuscript. The study was completed under government theme AAAA–A16–116021660078–0 with financial support from the Russian Foundation for Basic Research (grants RFBR Nos. 15–04–02029a, 15–29–02771) to NAP. Specimen preservation and examination were completed with financial support from the Russian Science Foundation (RSF grant No. 14–50–00029). SNN would like to thank Mr. Thi, Hanh, Phuong, Trang, Lam, Nong, Hieu for their permission and organization of fieldwork in Hon Ba NR and thank The Rufford Foundation for financial support. JC was supported by the programs of the National Natural Science Foundation of China (31622052), Southeast Asia Biodiversity Research Institute, Chinese Academy of Sciences (Y4ZK111B01: 2015CASEABRI002), the Youth Innovation Promotion Association CAS, and the Animal Branch of the Germplasm Bank of Wild Species, Chinese Academy of Sciences (the Large Research Infrastructure Funding). SM was supported by a Field Museum of Natural History Science Visiting Scholarship, for FMNH specimen study, and would like to thank the following for providing access to specimens and tissues: Barry Clark and Jeffrey Streicher (NHMUK), Darrel Frost and David Kizirian (AMNH), Julie Feinstein (AMCC), and Alan Resetar (FMNH). SM thanks Emma Teeling for laboratory support and resources.

References

- Bain RH, Nguyen TQ, Doan KV (2007) New herpetofaunal records from Vietnam. *Herpetological Review* 38: 107–117.
- Bonaparte CLJL (1850) *Conspectus Systematum. Herpetologiae et Amphibiologiae*. Editio altera reformata. Brill EJ, Lugdini Batavorum, 8 pp. [In Latin]
- Boulenger GA (1903) Descriptions of three new batrachians from Tonkin. *Annals and Magazine of Natural History*. Series 7 12: 186–188. <https://doi.org/10.1080/00222930308678835>
- Bourret R (1937) Notes herpétologiques sur l'Indochine française. XIV. Les batraciens de la collection du Laboratoire des Sciences Naturelles de l'Université. Descriptions de quinze espèces ou variétés nouvelles. Annexe au Bulletin Général de l'Instruction Publique. Hanoi 1937: 5–56. [In French]
- Bourret R (1942) Les Batraciens de l'Indochine. Institut Océanographique de l'Indochine, Hanoi, 453 pp. [In French]
- Chan KO, Blackburn DC, Murphy RW, Stuart BL, Emmett DA, Ho CT, Brown RM (2013) A new species of narrow-mouthed frog of the genus *Kaloula* from eastern Indochina. *Herpetologica* 69: 329–341. <https://doi.org/10.1655/HERPETOLOGICA-D-12-00094>
- Chan-ard T (2003) *A Photographic Guide to Amphibians in Thailand*. Darnsutha Press Co., Ltd., Bangkok, Thailand, 176 pp. [In Thai]
- Chen JM, Zhou WW, Poyarkov Jr. NA, Stuart BL, Brown RM, Lathrop A, Wang YY, Yuan ZY, Jiang K, Hou M, Chen HM, Suwannapoom C, Nguyen NS, Duong VT, Papenfuss TJ, Murphy RW, Zhang YP, Che J (2017) A novel multilocus phylogenetic estimation reveals unrecognized diversity in Asian horned toads, genus *Megophrys* sensu lato (Anura: Megophryidae). *Molecular Phylogenetics and Evolution* 106: 28–43. <https://doi.org/10.1016/j.ympev.2016.09.004>
- De Koninck R (1999) *Deforestation in Viet Nam*. International Research Centre, Ottawa, 101 pp.
- Delorme M, Dubois A, Grosjean S, Ohler A (2006) Une nouvelle ergotaxinomie des Megophryidae (Amphibia, Anura). *Alytes*, Paris 24: 6–21. [In French]
- Dubois A (1980) Notes sur la systematique et la répartition des amphibiens anoures de Chine et des régions avoisinantes IV. Classification générique et subgénérique des Pelobatidae Megophryinae. *Bulletin Mensuel de la Société Linnéenne de Lyon* 49: 469–482. <https://doi.org/10.3406/linly.1980.10444> [In French]
- Dubois A (1987) “1986” *Miscellanea taxinomica batrachologica* (I). *Alytes*, Paris 5: 7–95.
- Fei L, Ye C-y, Huang Y-a, Liu M-y (1999) [Atlas of Amphibians of China]. Henan Press of Science and Technology, 432 pp. [In Chinese]
- Fei L, Ye C-y, Huang Y-z, Jiang J-p, Xie F (2005) *An Illustrated Key to Chinese Amphibians*. Sichuan Publishing House of Science and Technology, Chongqing, 341 pp. [In Chinese]
- Fei L, Hu S-q, Ye C-y, Huang Y-z (2009) *Fauna Sinica. Amphibia. Volume 2. Anura*. Chinese Academy of Science; Science Press, Beijing, 957 pp. [In Chinese]
- Fei L, Ye C-y, Jiang J-p (2010) *Colored Atlas of Chinese Amphibians*. Sichuan Publishing Group/Sichuan Publishing House of Science and Technology, 519 pp. [In Chinese]
- Fei L, Ye C-y, Jiang J-p (2012) *Colored Atlas of Chinese Amphibians and Their Distributions*. Sichuan Publishing House of Science and Technology, Sichuan, China, 619 pp. [In Chinese]

- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39(4): 783–791. <https://doi.org/10.2307/2408678>
- Frost DR [Ed.] (1985) *Amphibian Species of the World. A Taxonomic and Geographical Reference*. Association of Systematics Collections and Allen Press, Lawrence, Kansas, USA, 732 pp.
- Frost DR, Grant T, Faivovich J, Bain RH, Haas A, Haddad CFB, de Sá RO, Channing A, Wilkinson M, Donnellan SC, Raxworthy CJ, Campbell JA, Blotto BL, Moler PE, Drewes RC, Nussbaum RA, Lynch JD, Green DM, Wheeler WC (2006) The amphibian tree of life. *Bulletin of the American Museum of Natural History* 297: 1–370. [https://doi.org/10.1206/0003-0090\(2006\)297\[0001:TATOL\]2.0.CO;2](https://doi.org/10.1206/0003-0090(2006)297[0001:TATOL]2.0.CO;2)
- Geissler G, Poyarkov Jr. NA, Grismer LL, Nguyen TQ, An HT, Neang T, Kupfer A, Ziegler T, Böhme W, Müller H (2014) New *Ichthyophis* species from Indochina (Gymnophiona, Ichthyophiidae): 1. The unstriped forms with descriptions of three new species and the redescrptions of *I. acuminatus* Taylor, 1960, *I. youngorum* Taylor, 1960 and *I. laosensis* Taylor, 1969. *Organisms Diversity and Evolution*, 1–32. <https://doi.org/10.1007/s13127-014-0190-6>
- Geissler P, Hartmann T, Ihlow F, Rödder D, Poyarkov Jr. NA, Nguyen TQ, Ziegler T, Böhme W (2015) The Lower Mekong: an insurmountable barrier for amphibians in southern Indochina. *Biological Journal of the Linnean Society* 114(4): 905–914. <https://doi.org/10.1111/bij.12444>
- Gerhardt HC, Huber F (2002) *Acoustic Communication in Insects and Anurans: Common Problems and Diverse Solutions*. University of Chicago Press, Chicago, 531 pp.
- Gorham SW (1974) Checklist of World Amphibians up to January 1, 1970. Saint-John, The New Brunswick Museum, 173 pp.
- Gosner KL (1960) A simplified table for staging Anura embryos and larvae with notes on identification. *Herpetologica* 16: 183–190.
- Goutte S, Dubois A, Howard SD, Marquez R, Rowley JJ, Dehling JM, Grandcolas P, Rongchuan X, Legendre F (2016) Environmental constraints and call evolution in torrent-dwelling frogs. *Evolution* 70(4): 811–826. <https://doi.org/10.1111/evo.12903>
- Gray JE (1825) A synopsis of the genera of reptiles and Amphibia, with description of some new species. *Annals of Philosophy. Series 2* 10: 193–217.
- Grosjean S (2001) The tadpole of *Leptobrachium (Vibrissaphora) echinatum* (Amphibia, Anura, Megophryidae). *Zoosystema* 23: 143–156.
- Grosjean S (2003) A redescription of the external and buccopharyngeal morphology of the tadpole of *Ophryophryne microstoma* Boulenger, 1903 (Megophryidae). *Alytes* 21(1–2): 45–58.
- Guibé J (1950) Catalogue des types d'Amphibiens du Muséum National d'Histoire Naturelle. Paris, 71 pp. [In French]
- Günther A (1864) *The Reptiles of British India*. The Ray Society, London, 452 pp + I–XXVII + 1–26 pl.
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nuclear Acids Symposium. Series 41*: 95–98.
- Hamidy A, Matsui M, Nishikawa K, Belabut DM (2012) Detection of cryptic taxa in *Leptobrachium nigrops* (Amphibia, Anura, Megophryidae), with description of two new species. *Zootaxa* 3398: 22–39.

- Hartmann T, Geissler P, Poyarkov Jr. NA., Ihlow F, Galoyan EA, Rödder D, Böhme W (2013) A new species of the genus *Calotes* Cuvier, 1817 (Squamata: Agamidae) from southern Vietnam. *Zootaxa* 3599(3): 246–260. <https://doi.org/10.11646/zootaxa.3599.3.3>
- Hedges SB (1994) Molecular evidence for the origin of birds. *Proceedings of the National Academy of Sciences of the United States of America* 91: 2621–2624.
- Hillis DM, Moritz C, Mable BK (1996) *Molecular Systematics*. 2nd edition. Sinauer Associates, Inc., Sunderland, Massachusetts, xvi + 655 pp.
- Huang Y, Fei L, Ye C (1991) Studies on internal oral structures of tadpoles of Chinese Pelobatidae. *Acta Biologica Sinica* 10: 71–99. [In Chinese]
- Huelsenbeck JP, Hillis DM (1993) Success of phylogenetic methods in the four-taxon case. *Systematic Biology* 42: 247–264. <https://doi.org/10.1093/sysbio/42.3.247>
- Huelsenbeck JP, Ronquist F (2001) MRBAYES: Bayesian inference of phylogeny. *Bioinformatics* 17: 754–755. <https://doi.org/10.1093/bioinformatics/17.8.754>
- Inger RF, Orlov NL, Darevsky IS (1999) *Frogs of Vietnam: A report on new collections*. Fieldiana. Zoology. New Series 92: 1–46.
- IUCN Standards and Petitions Subcommittee (2016) *Guidelines for Using the IUCN Red List Categories and Criteria*. Version 12. Prepared by the Standards and Petitions Subcommittee, 101 pp. <http://www.iucnredlist.org/documents/RedListGuidelines.pdf>
- Jiang K, Yan F, Suwannapoom C, Chomdej S, Che J (2013) A new species of the genus *Leptolalax* (Anura: Megophryidae) from northern Thailand. *Asian Herpetological Research* 4: 100–108. <https://doi.org/10.3724/SP.J.1245.2013.00100>
- Jobb G, Von Haeseler A, Strimmer K (2004) TREEFINDER: a powerful graphical analysis environment for molecular phylogenetics. *BMC Evolutionary Biology* 4: 18. <https://doi.org/10.1186/1471-2148-4-18>
- Khonsue W, Thirakhupt K (2001) A checklist of the amphibians of Thailand. *Natural History Journal of Chulalongkorn University* 1: 69–82.
- Kou Z-t (1985) A new species of *Ophryophryne* from Yunnan. *Acta Herpetologica Sinica/ Liangqi baxing dongwu yanjiu*. New Series, Chengdu 4(1): 41–43. [In Chinese]
- Kuhl H, Van Hasselt JC (1822a) *Uittreksels uit breieven van de Heeren Kuhl en van Hasselt, aan de Heeren C. J. Temminck, Th. van Swinderen en W. de Haan*. *Algemeene Konst-en Letter-Bode* 7: 99–104. [In Dutch]
- Kuznetsov AN, Kuznetsova SP (2011) Chuong 2. Thực Vật VQG Bidoup–Núi Bà [Chapter 5. Reptiles and Amphibians of Bidoup–Nui Ba National Park]. In: Nguyen DH, Kuznetsov AN (Eds) *Da Dang Sinh Hoc Va Dac Trung Sinh Thai Vuon Quoc Gia Bidoup–Nui Ba*, Nha Xuat Ban Hoa Hoc Tu Nhien Va Cong Nghe Publishing House, Hanoi, 37–105. [In Vietnamese]
- Laurance WF (2007) Forest destruction in tropical Asia. *Current Science* 93(11): 1544–1550.
- Liu C-c, Hu S-q (1962) A herpetological report of Kwangsi. *Acta Zoologica Sinica/ Dong wu xue bao*, Beijing 14(Supplement): 73–104. [In Chinese]
- Mahony S (2011a) Two new species of *Megophrys* Kuhl and Van Hasselt (Amphibia: Megophryidae), from western Thailand and southern Cambodia. *Zootaxa* 2734: 23–39.
- Mahony S (2011b) Erratum: Mahony, Stephen (2011) Two new species of *Megophrys* Kuhl and Van Hasselt (Amphibia: Megophryidae), from western Thailand and southern Cambodia. *Zootaxa* 2734: 23–39. *Zootaxa* 2924: 68.

- Mahony S, Teeling EC, Biju SD (2013) Three new species of horned frogs, *Megophrys* (Amphibia: Megophryidae), from northeast India, with a resolution to the identity of *Megophrys boettgeri* populations reported from the region. *Zootaxa* 3722(2): 143–169.
- Mahony S, Foley NM, Biju SD, Teeling EC (2017) Evolutionary history of the Asian Horned Frogs (Megophryinae): integrative approaches to timetree dating in the absence of a fossil record. *Molecular Biology and Evolution* 34(3): 744–771. <https://doi.org/10.1093/molbev/msw267>
- Manthey U, Grossmann W (1997) Amphibien und Reptilien Südasiens. Natur und Tier, Münster, 512 pp. [In German]
- Matsui M, Shimada T, Liu WZ, Maryati M, Khonsue W, Orlov N (2006) Phylogenetic relationships of Oriental torrent frogs in the genus *Amolops* and its allies (Amphibia, Anura, Ranidae). *Molecular Phylogenetics and Evolution* 38: 659–666. <https://doi.org/10.1016/j.ympev.2005.11.019>
- Matsui M, Hamidy A, Murphy RW, Khonsue W, Yambun P, Shimada T, Norhayati A, Daicus BM, Jiang JP (2010) Phylogenetic relationships of megophryid frogs of the genus *Leptobrachium* (Amphibia, Anura) as revealed by mtDNA gene sequences. *Molecular Phylogenetics and Evolution* 56: 259–272. <https://doi.org/10.1016/j.ympev.2010.03.014>
- Matsui M, Nishikawa K, Yambun P (2014a) A new *Leptolalax* from the mountains of Sabah, Borneo (Amphibia, Anura, Megophryidae). *Zootaxa* 3753: 440–452. <https://doi.org/10.11646/zootaxa.3753.5.3>
- Meijer W (1973) Devastation and Regeneration of Lowland Dipterocarp Forests in Southeast Asia. *BioScience* 23(9): 528–533. <https://doi.org/10.2307/1296481>
- Meyfroidt P, Lambin EF (2008) Forest transition in Vietnam and its environmental impacts. *Global Change Biology* 14(6): 1319–1336. <https://doi.org/10.1111/j.1365-2486.2008.01575.x>
- Murphy WJ, O'Brien SJ (2007) Designing and optimizing comparative anchor primers for comparative gene mapping and phylogenetic inference. *Nature Protocols* 2(11): 3022–3030. <https://doi.org/10.1038/nprot.2007.429>
- Nazarov R, Poyarkov NA, Orlov NL, Phung TM, Nguyen TT, Hoang DM, Ziegler T (2012) Two new cryptic species of the *Cyrtodactylus irregularis* complex (Squamata: Gekkonidae) from southern Vietnam. *Zootaxa* 3302: 1–24.
- Nguyen TQ, Phung TM, Schneider N, Botov A, Tran DTA, Ziegler T (2014) New records of amphibians and reptiles from southern Vietnam. *Bonn zoological Bulletin* 63(2): 148–156.
- Nguyen SV, Ho CT, Nguyen TQ (2005) Danh Lục Ech Nhái Và Bò Sát Việt Nam [A Checklist of the Amphibians and Reptiles of Vietnam]. Hanoi: Nhà Xuất Bản Hồng Nghiep, 180 pp. [In Vietnamese]
- Nguyen VS, Ho TC (1996) Danh lục bò sát và êch nhái Việt Nam. Hà Nội, Nhà xuất bản khoa học và kỹ thuật: 1–264. [In Vietnamese]
- Nguyen VS, Ho TC, Nguyen QT (2009) Herpetofauna of Vietnam. Edition Chimaira, Frankfurt am Main, 768 pp.
- Noble GK (1926) An analysis of the remarkable cases of distribution among the Amphibia, with descriptions of new genera. *American Museum Novitates* 212: 1–24.
- Ohler A (2003) Revision of the genus *Ophryophryne* Boulenger, 1903 (Megophryidae) with description of two new species. *Alytes*. Paris 21: 23–42.

- Ohler A, Wollenberg KC, Grosjean S, Hendrix R, Vences M, Ziegler T, Dubois A (2011) Sorting out *Lalos*: description of new species and additional taxonomic data on megophryid frogs from northern Indochina (genus *Leptolalax*, Megophryidae, Anura). *Zootaxa* 3147: 1–83.
- Orlov NL, Poyarkov Jr. NA., Nguyen TT (2015) Taxonomic notes on *Megophrys* frogs (Megophryidae: Anura) of Vietnam, with description of a new species. *Russian Journal of Herpetology* 22(3): 206–218.
- Orlov NL, Poyarkov Jr. NA., Vassilieva AB, Ananjeva NB, Nguyen TT, Sang NV, Geissler P (2012) Taxonomic notes on rhacophorid frogs (Rhacophorinae: Rhacophoridae: Anura) of southern part of Annamite Mountains (Truong Son, Vietnam), with description of three new species. *Russian Journal of Herpetology* 19: 23–64.
- Orlov NL, Murphy RW, Ananjeva NB, Ryabov SA, Ho CT (2002) Herpetofauna of Vietnam, a checklist. Part 1. Amphibia. *Russian Journal of Herpetology* 9: 81–104.
- Orlov NL, Nguyen SN, Ho CT (2008) Description of a new species and new records of *Rhacophorus* genus (Amphibia: Anura: Rhacophoridae) with the review of amphibians and reptiles diversity of Chu Yang Sin National Park (Dac Lac Province, Vietnam). *Russian Journal of Herpetology* 15: 67–84.
- Orlov NL, Ananjeva NB (2007) Amphibians of South-East Asia. St. Petersburg University Press, St. Petersburg, 270 pp. [In Russian]
- Orlov NL, Ho CT (2005) A new species of *Philautus* from Vietnam (Anura: Rhacophoridae). *Russian Journal of Herpetology* 12: 135–142.
- Palumbi SR (1996) Nucleic acids II: the polymerase chain reaction. In: Hillis DM, Moritz C, Mable BK (Eds) *Molecular systematics*. Second Edition. Sinauer Associates, Sunderland, 205–247.
- Penna M, Pottstock H, Velasquez N (2005) Effect of natural and synthetic noise on evoked vocal responses in a frog of the temperate austral forest. *Animal Behaviour* 70(3): 639–651. <https://doi.org/10.1016/j.anbehav.2004.11.022>
- Posada D, Crandall KA (1998) MODELTEST: testing the model of DNA substitution. *Bioinformatics* 14(9): 817–818. <https://doi.org/10.1093/bioinformatics/14.9.817>
- Poyarkov Jr. NA., Rowley JJJ, Gogoleva SS, Vassilieva AB, Galoyan EA, Orlov NL (2015a) A new species of *Leptolalax* (Anura: Megophryidae) from the western Langbian Plateau, southern Vietnam. *Zootaxa* 3931(2): 221–252. <https://doi.org/10.11646/zootaxa.3931.2.3>
- Poyarkov Jr. NA., Orlov NL, Moiseeva AV, Pawangkhanant P, Ruangsuwan T, Vassilieva AB, Galoyan EA, Nguyen TT, Gogoleva SS (2015b) Sorting out moss frogs: mtDNA data on taxonomic diversity and phylogenetic relationships of the Indochinese species of the genus *Theloderma* (Anura, Rhacophoridae). *Russian Journal of Herpetology* 22(4): 241–280.
- Poyarkov Jr. NA., Che J, Min M-S, Kuro-o M, Yan F, Li Ch, Iizuka K, Vieites DR (2012) Review of the systematics, morphology and distribution of Asian Clawed Salamanders, genus *Onychodactylus* (Amphibia, Caudata: Hynobiidae), with the description of four new species. *Zootaxa* 3465: 1–106.
- Poyarkov Jr. NA., Vassilieva AB, Orlov NL, Galoyan EA, Tran TAD, Le DTT, Kretova VD, Geissler P (2014) Taxonomy and distribution of narrow-mouth frogs of the genus *Microhyla* Tschudi, 1838 (Anura: Microhylidae) from Vietnam with descriptions of five new species. *Russian Journal of Herpetology* 21(2): 89–148.

- Poyarkov [Paiarkov] NA, Vasilieva AB (2011) Chuong 5. Bò Sát — Lường củ VQG (Vườn Quốc Gia) Bidoup–Núi Bà [Chapter 5. Reptiles and Amphibians of Bidoup–Núi Ba National Park]. In: Nguyen DH, Kuznetsov AN (Eds) Đa Dạng Sinh Học Và Đặc Trưng Sinh Thái Vườn Quốc Gia Bidoup–Núi Bà, Hanoi, Nhà Xuất Bản Khoa Học Tự Nhiên Và Công Nghệ Publishing House, pp 169–220. [In Vietnamese]
- Preininger D, Böckle M, Hödl W (2007) Comparison of anuran acoustic communities of two habitat types in the Danum Valley Conservation Area, Sabah, Malaysia. *Salamandra* 43(3): 129–138.
- Puillandre N, Lambert A, Brouillet S, Achaz G (2012) ABGD, Automatic Barcode Gap Discovery for primary species delimitation. *Molecular Ecology* 21: 1864–1877. <https://doi.org/10.1111/j.1365-294X.2011.05239.x>
- Pyron RA, Wiens JJ (2011) A large-scale phylogeny of Amphibia including over 2800 species, and a revised classification of advanced frogs, salamanders, and caecilians. *Molecular Phylogenetics and Evolution* 61: 543–583. <https://doi.org/10.1016/j.ympev.2011.06.012>
- Rao D-q, Yang D-t (1997) The karyotypes of Megophryinae (Pelobatinae) with a discussion on their classification and phylogenetic relationships. *Asiatic Herpetological Research* 7: 93–102. <https://doi.org/10.5962/bhl.part.18858>
- Ronquist F, Huelsenbeck JP (2003) MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19: 1572–1574. <https://doi.org/10.1093/bioinformatics/btg180>
- Rowley JJJ, Hoang DH, Le TTD, Dau QV, Cao TT (2010a) A new species of *Leptolalax* (Anura: Megophryidae) from Vietnam and further information on *Leptolalax tuberosus*. *Zootaxa* 2660: 33–45.
- Rowley J, Brown R, Kusirini M, Inger R, Stuart B, Wogan G, Chan-ard T, Cao TT, Diesmos A, Iskandar DT, Lau M, Ming LT, Makchai S, Neang T, Nguyen QT, Phimmachak S (2010b) Impending conservation crisis for Southeast Asian amphibians. *Biology Letters* 6: 336–338. <https://doi.org/10.1098/rsbl.2009.0793>
- Rowley JJJ, Le TTD, Tran TAD, Stuart BL, Hoang DH (2010c) A new tree frog of the genus *Rhacophorus* (Anura: Rhacophoridae) from southern Vietnam. *Zootaxa* 2727: 45–55.
- Rowley JJJ, Le DTT, Tran DTA, Hoang DH (2011a) A new species of *Leptolalax* (Anura: Megophryidae) from southern Vietnam. *Zootaxa* 2796: 15–28.
- Rowley JJJ, Le TTD, Hoang DH, Dau QV, Cao TT (2011b) Two new species of *Theloderma* (Anura: Rhacophoridae) from Vietnam. *Zootaxa* 3098: 1–20.
- Rowley JJJ, Hoang HD, Dau VQ, Le TTD (2012) A new species of *Leptolalax* (Anura: Megophryidae) from central Vietnam. *Zootaxa* 3321: 56–68.
- Rowley JJJ, Tran DTA, Le DTT, Dau VQ, Peloso PLV, Nguyen TQ, Hoang HD, Nguyen TT, Ziegler T (2016) Five new, microendemic Asian Leaf-litter Frogs (*Leptolalax*) from the southern Annamite mountains, Vietnam. *Zootaxa* 4085: 63–102. <https://doi.org/10.11646/zootaxa.4085.1.3>
- Schneider JG (1799) *Historia Amphibiorum Naturalis et Literariae*. Fasciculus Primus. Continentes Ranas, Calamitas, Bufones, Salamandras et Hydros in Genera et Species Descriptos Notisque usis Distinctos. Friederici Frommanni, Jena, 304 pp.
- Shimada T, Matsui M, Yambun P, Sudin A (2011) A taxonomic study of Whitehead's torrent frog, *Meristogenys whiteheadi*, with descriptions of two new species (Amphibia: Ranidae).

- Zoological Journal of the Linnean Society 161: 157–183. <https://doi.org/10.1111/j.1096-3642.2010.00641.x>
- Smith MA (1921) New or little-known reptiles and batrachians from southern Annam (Indo-China). Proceedings of the Zoological Society of London 1921: 423–440.
- Smith MA (1924) New tree-frogs from Indo-China and the Malay Peninsula. Proceedings of the Zoological Society of London 1924: 225–234. <https://doi.org/10.1111/j.1096-3642.1924.tb01499.x>
- Stuart BL (2005) New frog records from Laos. Herpetological Review 36: 473–479.
- Stuart BL, Rowley JJL, Neang T, Emmett DA, Sitha S (2010) Significant new records of amphibians and reptiles from Virachey National Park, northeastern Cambodia. Cambodian Journal of Natural History 2010: 38–47.
- Stuart BL, Sok K, Neang T (2006) A collection of amphibians and reptiles from hilly eastern Cambodia. Raffles Bulletin of Zoology 54: 129–155.
- Stuart BL, Phimmachak S, Seateun S, Sivongxay N (2012) A new *Leptobrachium* (Anura: Megophryidae) from the highlands of southeastern Laos. Zootaxa 3155: 29–37.
- Stuart BL, Rowley JJL, Tran DTA, Le DTT, Hoang DH (2011) The *Leptobrachium* (Anura: Megophryidae) of the Langbian Plateau, southern Vietnam, with description of a new species. Zootaxa 2804: 25–40.
- Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. Molecular Biology and Evolution 30: 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Taylor EH (1962) The amphibian fauna of Thailand. The University of Kansas Science Bulletin, XLIII, 255–599. <https://doi.org/10.5962/bhl.part.13347>
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. Nucleic Acids Research 25: 4876–4882. <https://doi.org/10.1093/nar/25.24.4876>
- Tian W, Hu Q (1983) Taxonomic study on genus *Megophrys*, with descriptions of two new genera. Acta Herpetologica Sinica. New Series 2(2): 41–48. [In Chinese]
- Tian W, Hu Q (1985) Taxonomical studies on the primitive anurans of the Hengduan mountains, with descriptions of a new subfamily and subdivision of *Bombina*. Acta herpetologica Sinica 4: 219–224. [In Chinese]
- Tschudi JJv (1838) Classification de Batrachier mit Berücksichtigung der fossilen Thiere dieser Abtheilung der Reptilien. Neuschâtel, Petitpierre.
- Vassilieva AB, Galoyan EA, Poyarkov NA, Geissler P (2016) A Photographic Field Guide to the Amphibians and Reptiles of the Lowland Monsoon Forests of Southern Vietnam. Frankfurt Contributions to Natural History, Vol. 36. Chimaira Buchhandelsgesellschaft mbH, Frankfurt-am-Main, 319 pp.
- Vassilieva AB, Galoyan EA, Gogleva SI, Poyarkov NA Jr (2014) Two new species of *Kalophrynus* Tschudi, 1838 (Anura: Microhylidae) from the Annamite mountains in southern Vietnam. Zootaxa 3796(3): 401–434. <https://doi.org/10.11646/zootaxa.3796.3.1>
- Vences M, Thomas M, Bonett RM, Vieites DR (2005a) Deciphering amphibian diversity through DNA barcoding: chances and challenges. Philosophical Transactions of the Royal Society London, Ser. B 360: 1859–1868.

- Vences M, Thomas M, Van der Meijden A, Chiari Y, Vieites DR (2005b) Comparative performance of the 16S rRNA gene in DNA barcoding of amphibians. *Frontiers in Zoology* 2: 5. <https://doi.org/10.1186/1742-9994-2-5>
- Vieites DR, Wollenberg KC, Andreone F, Kohler J, Glaw F, Vences M (2009) Vast underestimation of Madagascar's biodiversity evidenced by an integrative amphibian inventory. *Proceedings of the National Academy of Sciences of the United States of America* 106: 8267–8272. <https://doi.org/10.1073/pnas.0810821106>
- Wang Y, Zhang T-d, Zhao J, Sung Y-h, Yang J, Pang H, Zhang Z (2012) Description of a new species of the genus *Xenophrys* Günther, 1864 (Amphibia: Anura: Megophryidae) from Mount Jinggang, China, based on molecular and morphological data. *Zootaxa* 3546: 53–67.
- Wells KD (2007) *The Ecology and Behaviour of Amphibians*. Chicago University Press, 1148 pp. <https://doi.org/10.7208/chicago/9780226893334.001.0001>
- Wilkinson JA, Drewes RC, Tatum OL (2002) A molecular phylogenetic analysis of the family Rhacophoridae with an emphasis on the Asian and African genera. *Molecular Phylogenetics and Evolution* 24: 265–273. [https://doi.org/10.1016/S1055-7903\(02\)00212-9](https://doi.org/10.1016/S1055-7903(02)00212-9)
- Xu Q, Liu S, Wan R, Yue B, Zhang X (2014) The complete mitochondrial genome of the *Vibrissaphora boringii* (Anura: Megophryidae). *Mitochondrial DNA* 27(1): 758–759. <https://doi.org/10.3109/19401736.2014.915527>
- Yang D-t (Ed.) (1991) *Yunnan liang qi lei zhi* [Amphibian Fauna of Yunnan]. China Forestry Publishing House, Beijing, China, 259 pp. [In Chinese]
- Yang D-t (2008) Amphibia. In: Yang D-t, Rao D-q (Eds) *Amphibia and Reptilia of Yunnan*. Yunnan Publishing Group Corporation, Yunnan Science and Technology Press, Kunming, 12–152. [In Chinese]
- Ye C-y, Fei L, Hu S-q (1993) *Rare and Economic Amphibians of China*. Chengdu, China: Sichuan Publishing House of Science and Technology, 412 pp. [In Chinese]
- Zhang P, Liang D, Mao R-L, Hillis DM, Wake DB, Cannatella D (2013) Efficient Sequencing of Anuran mtDNAs and a Mitogenomic Exploration of the Phylogeny and Evolution of Frogs. *Molecular Biology and Evolution* 30(8): 1899–1915. <https://doi.org/10.1093/molbev/mst091>
- Zhang P, Papenfuss TJ, Wake MH, Qu L, Wake DB (2008) Phylogeny and biogeography of the family Salamandridae (Amphibia: Caudata) inferred from complete mitochondrial genomes. *Molecular Phylogenetics and Evolution* 49(2): 586–597. <https://doi.org/10.1016/j.ympev.2008.08.020>
- Zhang Y-x, Wen Y-t (2000) *Amphibians in Guangxi*. Guangxi Biodiversity Studies. Guilin, China, Guangxi Normal University Press, 183 pp. [In Chinese]
- Zheng Y, Zeng X, Yuan Y, Liu Z (2004) Phylogenetic positions of *Ophryophryne* and four *Leptobrachium* group genera in Megophryidae (Anura). *Sichuan Dong Wu* 23(3): 290–295. [In Chinese]

Appendix I

Examined material, museum IDs given in bold.

Ophryophryne elfina sp. n.: **ZMMU A-5669** (Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; NAP-02658; 1 adult male; holotype); **ZMMU A-5170** (Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Bidoup Mt.; ABV-00455; ABV-00454; ABV-00472; ABV-00471; 4 adults; paratypes); **ZMMU A-4788** (Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Hon Giao Mt., Bidoup Mt.; NAP-01455; NAP-01449; NAP-01450; NAP-01460; 4 adults; paratypes); **ZMMU A-5674** (Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Hon Giao Mt.; NAP-01451; NAP-01452; 2 adults; paratypes); **ZMMU A-5675** (Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Bidoup Mt.; NAP-01456; NAP-01459; 2 adults; paratypes); **ZMMU A-5691** (Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; ABV-00580, ABV-00581; 2 juveniles; paratypes); **ZMMU A-5650** (Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; 10 adults); **ZISP 12836–12879** (Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; 44 adults; tentative identification); **ZMMU A-5665** (Vietnam, Dak Lak, Chu Yang Sin N.P.; ROM 36348; ROM 36289; ROM 36554; ROM 36470; 4 adults); **ZISP 12880–12884** (Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; ROM 36471; ROM 36472; ROM 36469; ROM 36366; ROM 36349; 5 adults); **ZMMU A-3937** (Vietnam, Khanh Hoa Prov., Hon Ba N.R.; 1 adult); **ZMMU A-4935** (Vietnam, Khanh Hoa Prov., Hon Ba N.R.; ABV-00316; 1 adult); **ZMMU A-4716** (Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Hon Giao Mt., Bidoup Mt.; NAP-01871; NAP-01782; NAP-01758; NAP-01757; NAP-01783; 5 adults); **ITBCZ 2786, 2788, 2792, 2828, 2908–09, 2918–19, 3502** (Vietnam, Khanh Hoa Prov., Hon Ba N.R.; 9 adults); **ZMMU A-5679** (Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Bidoup Mt.; NAP-01169; 7 larvae); **ZMMU A-5684** (Vietnam, Dak Lak Prov., Chu Yang Sin N.P., Chu Pan Fan Mt.; NAP-02673; 4 larvae).

Ophryophryne gerti: **BMNH 1921.4.1.324** (Vietnam, Lam Dong Prov., south-east of Da Lat, Cam Ly river; 1 adult male; holotype); **BMNH 1921.4.1.323** (Vietnam, Lam Dong Prov., Dran; 1 immature female; paratype); **ZMMU A-4714** (Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Giang Ly; NAP-01878; 1 adult); **ZMMU A-4715** (Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Giang Ly; NAP-01790; NAP-01788; 2 adults); **ZMMU A-4718** (Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Giang Ly; NAP-01789; 1 adult); **ZMMU A-4843** (Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; ABV-00067; 1 adult); **ZMMU A-5670** (Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; NAP-02758; NAP-02759; NAP-02760; NAP-02761; 4 adults); **ZMMU A-5673** (Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; ABV-00530; ABV-00577; 2 adults); **ZISP 12740–12747** (Vietnam, Dak Lak Prov., Chu Yang Sin N.P.; ROM 36522–36529; 8 adults).

Ophryophryne hansii: **FMNH 252879, FMNH 252880, FMNH 252882, FMNH 252884** (Vietnam, Gia-Lai Prov., Ankhe Dist., 60 km to the northwest from Kannack; 1 adult male; holotype; 2 adult males and 1 adult female; paratypes); **FMNH 252873** (Vietnam, Gia-Lai Prov., Ankhe Dist., 20 km to the northwest from Kan-

- nack; 1 adult male); **AMNH 163680** (Vietnam, Quang Nam Prov., Tra My Dist., Tra Don Commune; 1 adult male); **AMNH 169284** (Vietnam, Thua Tien Hue Prov., A Luoi Dist., Tram Tra Ve (Forestry station of Huong Giang State Forestry Enterprise); 1 adult male); **AMNH 169286** (Vietnam, Thua Tien Hue Prov., Huong Thuy Dist., Khe Dau Station; 1 adult male); **AMNH 161353** (Vietnam, Ha Tinh Prov., Huong Son, Huong Son Reserve; 1 adult male); **FMNH 258008–258009, FMNH 258046–258051** (Lao P.D.R., Xe Kong Prov., Kaleum Dist., Xe Sap National Biodiversity Conservation Area; 5 adult males and 4 adult females).
- Ophryophryne miscrostoma*: **BMNH 1947.2.22.50, BMNH 1947.2.22.52** (Vietnam, Mau Son Mts.; 2 adult females; lectotype & paralectotype); **AMNH 168682** (Vietnam, Lao Cai Prov., Van Ban Dist., Nam Tha Commune; 1 adult male).
- Ophryophryne pachyproctus*: **YU A8311032** (China, Yunnan Prov., Xishuangbanna Pref., Mengla Co., Yiwu, Zhoushihe river; 1 adult male; holotype); **YU A8311033–A8311037, YU A845099–A845100** (China, Yunnan Prov., Xishuangbanna Pref., Mengla Co., Yiwu, Zhoushihe river; 8 adult males; paratypes).
- Ophryophryne* cf. *poilani*: **AMNH 169287** (Vietnam, Thua Tien Hue Prov., A Luoi Dist., A Pat Forestry Protection Department Range Station; 1 adult female); **AMNH 163668** (Vietnam, Quang Nam Prov., Tra My Dist., Tra Don Commune; 1 adult female).
- Ophryophryne synoria*: **FMNH 262779** (Cambodia, Mondolkiri Prov., O’Rang Dist., O Chung Chry stream; 1 adult male; holotype); **FMNH 262778** (Cambodia, Mondolkiri Prov., O’Rang Dist., O Chung Chry stream; 1 adult male; paratype); **ZMMU A-5003** (Vietnam, Binh Phuoc Prov., Bu Gia Map N.P.; ABV-00379; ABV-00380; ABV-00376; ABV-00381; ABV-00377; 5 adults); **ZMMU A-4516** (Vietnam, Binh Phuoc Prov., Bu Gia Map N.P.; NAP-00729; NAP-00727; NAP-00728; NAP-00730; NAP-00731; 5 adults); **ZMMU A-4864** (Vietnam, Lam Dong Prov., Loc Bac forestry; ABV-00159; ABV-00209; 2 adults); **ZMMU A-5676** (Vietnam, Lam Dong Prov., Bidoup–Nui Ba N.P., Giang Ly; NAP-01438; NAP-01437; 2 adults).
- Ophryophryne* sp.: **BMNH 1972.1524** (Lao P.D.R., “Pak Maat”; 1 adult male; paratype of *O. gerti*).

Appendix 2

Factor coordinates of the morphometric characters used in PCA analysis, based on correlations (factors 1 to 3).

Character	factor 1	factor 2	factor 3
SVL	-0.969013	-0.052119	-0.061729
HW	-0.972214	-0.094198	-0.099264
HL	-0.963376	-0.112316	-0.089300
ED	-0.872338	-0.060587	0.293135
TYD	-0.853903	-0.302433	0.054377
TYE	-0.869868	-0.053510	-0.256344
SL	-0.817708	0.454058	-0.126722
EN	-0.774425	-0.018511	-0.529049
NS	-0.302259	0.915825	0.017344
IUE	-0.866068	-0.171158	0.204049
IN	-0.818811	0.248915	0.129748
UEW	-0.848819	0.181381	0.241783
FAL	-0.937173	-0.162120	-0.073309
HAL	-0.957549	0.054374	-0.001476
FIL	-0.931528	-0.047484	0.090017
FIIL	-0.939026	-0.049085	0.087852
FIIL	-0.963650	0.030269	-0.041650
FIVL	-0.902097	0.102388	-0.091818
SHL	-0.950468	-0.104557	0.046962
TL	-0.921529	-0.091021	0.019328
FOL	-0.918076	0.080433	0.100931
TFOL	-0.971401	0.003596	0.002474
IMT	-0.761261	-0.079858	0.068715

Appendix 3

Measurements of advertisement call temporal parameters and one-way ANOVA/Kruskal-Wallis results for comparison (* $p < 0.001$) between April and February sets of call recordings for *Ophryophryne elfina* sp. n. Seconds (s), milliseconds (ms).

Parameters	<i>O. elfina</i> sp. n. 10–15 April 2014	<i>O. elfina</i> sp. n. 10 February 2015	Tukey/ Mann- Whitney U post hoc tests	ANOVA/ Kruskal-Wallis results
Temperature of recording	11.3–11.4°C	17.0–17.5°C	–	–
Number of males	2	1	–	–
Number of recordings	3	2	–	–
Number of series	93	47	–	–
Number of calls	1301	496	–	–
Call repetition rate per recording (calls/s)	1.25 ± 0.35 (0.77–1.95) <i>N</i> = 3	1.07 ± 0.06 (1.01–1.12) <i>N</i> = 2	$p = 0.71$	$F_{1,3} = 0.2$
Number of calls per series	14 ± 0.45 (2–22) <i>N</i> = 93	10.53 ± 0.72 (3–21) <i>N</i> = 47	$p < 0.001$	$F_{1,138} = 18.2^*$
Series duration (s)	3.37 ± 0.1 (0.43– 5.05) <i>N</i> = 93	3.52 ± 0.27 (0.61–9) <i>N</i> = 47	$p = 0.52$	$F_{1,138} = 0.4$
Call repetition rate per series (calls/s)	4.22 ± 0.08 (2.92–5.49) <i>N</i> = 93	3.18 ± 0.1 (1.33–4.91) <i>N</i> = 47	$p < 0.001$	$F_{1,138} = 63.9^*$
Call duration (ms)	70 ± 0.27 (25–112) <i>N</i> = 1301	79 ± 0.32 (48–102) <i>N</i> = 496	$p < 0.001$ (<i>U</i> = 131097)	$H_{1,1798} = 380.2^*$
Inter-calls interval (ms)	184 ± 1.72 (96–621) <i>N</i> = 1209	271.17 ± 4.92 (102–942) <i>N</i> = 448	$p < 0.001$ (<i>U</i> = 109162)	$H_{1,1657} = 349.2^*$
Inter-series interval (s)	6.72 ± 0.54 (1.26–31.65) <i>N</i> = 90	6.09 ± 0.56 (1.42–22.14) <i>N</i> = 45	$p = 0.83$ (<i>U</i> = 2022)	$H_{1,136} = 0.05$